arXiv:math/0612001v4 [math.DG] 31 Jan 2009

A Real-Analytic Jordan Curve Cannot Bound Infinitely Many Relative Minima of Area

Michael Beeson
Department of Computer Science
San Jose State University
San Jose, California 95192
USA
email: beeson@cs.sjsu.edu

January 31, 2009

Abstract

Let Γ be a real-analytic Jordan curve in R^3. Then Γ cannot bound infinitely many disk-type minimal surfaces that provide relative minima of area.

1 Introduction

Branch points of minimal surfaces have been of interest ever since Douglas and Rado solved Plateau's problem in the 1930's, because their solution methods proved the existence of minimal surfaces which might in general have branch points. It was not until the seventies that the regularity (lack of branch points) of solutions providing an absolute minimum of area with given real-analytic boundary was proved. Readers not already intimately familiar with branch points may want to view the animated pictures posted to the Web at [5] for a visual introduction to branch points. One picture of a boundary branch point is given here in Figure 1.

Branch points of minimal surfaces not furnishing a relative minimum of area are also of considerable interest, because they are intimately connected with the "finiteness problem". It follows (as shown in [2], p. 121) from the work of Böhme [7] that if a real-analytic Jordan curve Γ bounds infinitely many disk-type relative minima of area, then it bounds an analytic one-parameter family of such minimal surfaces, all with the same Dirichlet's integral, which can be continued until it either loops or terminates in a branched minimal surface. In [27], it is proved that a loop of disk-type minimal surfaces contains one which is not a relative minimum of area. Therefore, to prove that Γ does not bound infinitely many relative minima of area, we only have to rule out the possibility that Γ bounds a one-parameter family of minimal surfaces $u(t, z)$, terminating in a branched minimal surface when the parameter t is zero, and furnishing a relative minimum of area for each positive t. Tomi solved the finiteness problem for absolute minima of area and real-analytic boundaries [27] in this way, since the limit of absolute minima of area is again an absolute minimum, and hence has no branch points. But the problem for relative minima has remained open. (A discussion of other open problems in this area is given at the end of the paper.)



Figure 1: This is a boundary branch point of order 2 and index 1. A boundary branch point of order $2m$ goes "around" $(2m+1)/2$ times while going "up and down" $2m+k+1)/2$ times. Could a one-parameter family of surfaces, each of which is a relative minima of area, and all with the same Jordan boundary, end in a surface like this?

Suppose we have a boundary branch point; orient the boundary so it is tangent to the X-axis with the branch point at origin. Consider the unit normal N restricted to the boundary. As the parameter t varies, we do not know *a priori* how N behaves, but at least it must remain perpendicular to the boundary. Close to the branch point it must almost lie in the YZ plane. Most of our paper is filled with calculations concerning the possible behavior of N as the parameter t goes to zero.

The finiteness problem for relative minima of area was first attacked in [2], where interior branch points arising as the limit of one-parameter families of relative minima were shown not to exist, and in [3], such boundary branch points were eliminated under special conditions on Γ, but not in general; so the finiteness problem for relative minima remained open until now, except for special boundaries.

The main line of attack in [2] is an eigenvalue argument, that in the case of an interior branch point, the Gaussian image of u for small positive t must include more than a hemisphere, and hence u cannot be a relative minimum of area. The reason why it must include more than a hemisphere is the Gauss-Bonnet-Sasaki-Nitsche formula, which expresses the total boundary curvature as a sum of contributions from the Gaussian area and the branch points. This formula implies that as soon as $t > 0$, the contribution of the branch point term to the total curvature when $t = 0$ must be made up by "extra" Gaussian area totaling $2m\pi$, where $2m$ is the order of the boundary branch point. For boundary branch points, this argument implies that for small positive t, there cannot be a neighborhood in the parameter domain on which the Gauss map covers more than the upper hemisphere. Near the boundary the normal is confined to within $O(t)$ of the YZ plane, and it seems plausible that the extra Gaussian area is contributed in the form of m hemispheres. However, that is difficult to prove. The behavior of N could, *a priori*, be quite complicated.

The main line of attack in [3] is the calculation of the eigenfunction corresponding to the first eigenvalue. The starting point of this calculation is the relation between the second variations of area and of Dirichlet's integral; namely, the condition for ϕ to belong to the kernel of the second variation of area is given by $\phi = k \cdot N$, where k belongs to the kernel of the second variation of area.[1] The "tangent vector" $k = u_t$ can be shown to be a "forced Jacobi direction" associated with the branch point; these are purely tangential, so as a function of the parameter t, the eigenfunction goes to zero. We calculate how the eigenfunction behaves on the w-plane, that is, on a region near the origin that shrinks to zero as t^γ. The leading term in t of this function is a harmonic function in the w-plane which inherits from the eigenfunction the property that it must have only one sign in the upper half plane. Using this property, we are able to derive additional information about the behavior of N near the origin.

In this paper, we go over the eigenfunction calculations again, for two reasons: First, in [3] we failed to include a term allowing for the possibility that there might be branch points in the lower half plane for $t > 0$ that might converge to the origin as t goes to zero. Second, we now simplify and improve the eigenfunction analysis.

We analyze the way in which the zeroes $a_i(t)$ of the function f in the Weierstrass representation approach zero as t goes to zero. These go to zero asymptotically in this way: $a_i(t) = \alpha_i t^\gamma$, for some γ. Possibly some of the a_i go to zero faster than others; we group the a_i into "rings", each ring containing the roots that go to zero with the same power of t. In the vicinity of (some of) these a_i the Gauss map of u^t for $t > 0$ must contribute enough Gaussian area to make about $2m\pi$ "extra" Gaussian area, which will disappear in the limit to account for the branch point term in the Gauss-Bonnet-Sasaki-Nitsche formula. Consider the behavior of the normal N on the boundary near the point where the branch point is when $t = 0$. The normal must be perpendicular to the boundary, so it is confined to the vicinity of the YZ-plane if we take Γ to be tangent to the X-axis. But its behavior there might be quite uncontrollable, and moreover, several of the a_i can be associated with the same α_i.

---

[1] Note that ϕ is a scalar, because with area it is enough to consider normal variations given by $u + \phi N$. On the other hand k is a vector since the second variation of Dirichlet's integral is defined as a Frechet derivative in the space of (vectors defining) surfaces u. The theorem that $\phi = k \cdot N$ is proved in [2].

In this argument there is another difficulty: the terms calculated for the eigenfunction might all cancel out, leaving the true eigenfunction hidden in the error terms. This is the case we call "H constant", since the eigenfunction is the imaginary part of H. It turns out that the main difficulty is to rule out this case. To do so, we first calculate exactly what the eigenfunction must look like in case H is constant. Then, we are able to show that if H is constant, no Gaussian area is contributed on the outermost "ring" of roots. We then "chase the Gaussian area" from one ring of roots down to the next, each time moving to a higher power of t and showing that if the terms in the eigenfunction cancel out to that power of t, then no Gaussian area can be contributed by the roots in that ring. But since *somewhere* we must pick up $2m\pi$ in extra Gaussian area, H cannot be constant "all the way down". Once that is proved, we can show that in fact it must not be constant on outermost ring, and all the roots go to zero with the same power of t.

At that point, the proof is half complete. We still must deal with the case when H is not constant. But our analysis of the eigenvalue and the behavior of the eigenfunction near the α_i tell us that each a_i is contributing exactly one hemisphere of Gaussian area. To finish the proof, we analyze the behavior of the eigenfunction for large w, where the surface is $u(z)$, and $z = t^\gamma w$. It turns out that the crucial issue here is whether $k/(2m+1)$ is an integer or not, where k is the index and $2m$ is the order of the branch point. If it is an integer, then the eigenfunction will have two signs in the upper half-plane, which is a contradiction. We are able to show that it must be an integer, by analyzing the behavior of the surface *on the boundary* near origin in two ways: once using the Weierstrass representation, and twice using the parametrization of the analytic boundary curve. Comparing the results of these two analyses we find that $k/(2m+1)$ is an integer, which completes the proof.

There is an appendix giving a "Dictionary of Notation", in which we list the symbols that are used in more than one section of the paper, and give a reminder of their definitions, as an aid to the reader.

We note in passing that as of yet, nobody has produced an example of a real-analytic Jordan curve bounding a minimal surface with a boundary branch point, although Gulliver has exhibited in [14] a surface with a C^∞ boundary and a boundary branch point. The experts I have asked all believe that they can exist, but an example is still missing. Also, it is not known whether Gulliver's example is area-minimizing or not.[2]

I am grateful to Daniel Wienholtz, Fritz Tomi, Tony Tromba, and Joel Spruck for fruitful discussions of the subject, and to Wienholtz and Tomi for their attention to my earlier work on this problem. Special thanks go to Fritz

---

[2] Wienholtz has used the solution of Björling's problem ([10], page 120) with boundary values $\Gamma(t^3)$ where Γ is a regular parametrization of a real-analytic arc, and a suitably specified normal, to produce examples of a minimal surface partially bounded by a real-analytic arc which is not a straight line segment, with a branch point on the arc.

Tomi for his careful reading of several preceding versions.

2 Preliminaries

By a real-analytic arc, or real-analytic Jordan curve, we mean an arc or curve that can be parametrized as a real-analytic function of arc length. We will consider surfaces bounded by a real-analytic Jordan curve Γ. We suppose that Γ passes through the origin tangent to the X-axis, and that u takes the portion of the real axis near origin onto Γ, with $u(0) = 0$. We still are free to orient the Y and Z axes. We do this in such a way that the normal at the branch point (which is well-defined) points in the positive Z-direction. With Γ oriented in this way, there will be two integers p and q such that Γ has a parametrization in the form

$$\Gamma(\tau) = \begin{bmatrix} \tau + O(\tau^2) \\ \frac{C_1\tau^{q+1}}{q+1} + O(\tau^{q+2}) \\ \frac{C_2\tau^{p+1}}{p+1} + O(\tau^{p+2}) \end{bmatrix}$$

for some nonzero real constants C_1 and C_2. But it is possible to choose τ more carefully so that we do not have the $O(\tau^2)$ term in the first coordinate. Let $\tau(z) = X(z) = \mathrm{Re}\frac{1}{2}\int f - fg^2 dz$. Then on the boundary we have

$$u(z) \quad = \quad \Gamma(\tau(z))$$

This parametrization and function $\tau(z)$ were inspired by Lewy's equation (see [11], p. 38).

We therefore have

$$\Gamma'(\tau) \quad = \quad \begin{bmatrix} 1 \\ C_1\tau^q + O(\tau^{q+1}) \\ C_2\tau^p + O(\tau^{p+1}) \end{bmatrix} \tag{1}$$

We write $u(z) = (X(z), Y(z), Z(z))$. We make use of the Enneper-Weierstrass representation of u (see e.g. [10], p. 108)

$$u(z) = Re \begin{bmatrix} \frac{1}{2}\int f - fg^2\,dz \\ \frac{i}{2}\int f + fg^2\,dz \\ \int fg\,dz \end{bmatrix}$$

where f is analytic and g is meromorphic in the upper half-disk.

Definition 1 *The* order *of the branch point is the order of the zero of f. The* index *of the branch point is the order of the zero of g.*

The order of a boundary branch point of a solution of Plateau's problem must be even (since the boundary is taken on monotonically). It is customary to write it as $2m$, and to use the letter k for the index. Thus $f(z) = z^{2m} + O(z^{2m+1})$ and $g(z) = cz^k + O(z^{k+1})$ for some constant c.

The Gauss-Bonnet formula says that for regular surfaces (minimal or not)

$$\int KW\,dx\,dy + \int_\Gamma \kappa_g = 2\pi$$

Note that for minimal surfaces, KW is negative. For minimal surfaces with branch points, there is another term in the Gauss-Bonnet formula:

$$\int KW\,dx\,dy + \int_\Gamma \kappa_g = 2\pi + 2M\pi$$

where M is the sum of the orders of the interior branch points and the half-orders of the boundary branch points. See [11], p. 121, or [22], p. 333.

Since the 1930s it has been known that, if Γ bounds infinitely many minimal surfaces, there is a sequence u_n of them converging to a minimal surface u bounded by Γ. We need an "analytic compactness" result that will give us a one-parameter family of minimal surfaces instead of just a sequence. Böhme [7] and Tromba [29] applied nonlinear global analysis to the theory of minimal surfaces. This is deep and beautiful work, and they (as well as Tomi) drew deep and beautiful consequences from it, such as [28], but we need only one result from it, the analytic compactness result of Böhme:

Theorem 1 (Böhme) *Let u be a minimal surface bounded by a real-analytic Jordan curve in R^3. Then either u is isolated, or there is a one-parameter family of minimal surfaces u^t defined for t in some interval; u^t is analytic in both variables simultaneously.*

This is not the most general theorem proved by Böhme. Others are stated in [2] and of course in [7], but this one suffices for our purposes.

The starting point of our work is the following theorem, derived from Böhme's theorem using the connections between the second variation of Dirichlet's integral and area, the connection between the second variation of area and a natural eigenvalue problem, the one-dimensionality of the eigenspace of the least eigenvalue, and an argument due to Tomi [27] ruling out "loops" in a one-parameter family of relative minima.

Theorem 2 *Let Γ be a real-analytic Jordan curve bounding infinitely many relative minimal of area. There there exists a one-parameter family of minimal surfaces u^t, bounded by Γ, defined for t in some interval containing 0, such that for $t > 0$, u^t is a relative minimum of area, and u^0 has an (interior or boundary) branch point.*

Proof. See [2], p. 121.

3 On the zero set of real-analytic functions

Several times in this paper we have reason to consider the zero set of a function that is analytic in two variables (one real and one complex). When we refer to a function of a real variable t and a complex variable z as "analytic", we mean it is given by a power series in z and t, so in particular it is real-analytic in t for fixed z and complex-analytic in z for fixed t. The following lemma describes the zero sets of such functions.

Lemma 1 *Let A be an open subset of $\mathbf{R} \times \mathbf{C}$ containing the point $(0, p)$. Let f be a complex-valued analytic function defined on A. Suppose $f(0, p) = 0$ and that for each sufficiently small $t > 0$, $f(t, \cdot)$ is not constant. Then there exists a neighborhood V of $(0, p)$ and finitely many functions c_i such that $f(t, c_i(t)) = 0$, every zero of f in V has the form $(t, c_i(t))$, and the $c_i(t)$ are analytic in some rational power of t, with $c_i(0) = p$.*

Proof. A classical theorem tells us that the zero set S is locally analytically triangulable. That is, there exist a neighborhood V of $(0, p)$ and finitely many closed simplexes and real-analytic homeomorphisms from those simplexes into S such that $V \cap S$ is contained in the range of those those simplexes. Six references for this theorem are given in [2], where the theorem is discussed on page 117. The theorem refers to *closed* simplexes, which means that the homeomorphisms in the conclusion are analytic even at the endpoints of intervals or boundary points of 2-simplexes.

If f does not depend on t at all, the theorem is trivial. Otherwise, if f is divisible by some power of t, we can divide that power out without changing the zero set of f, so we can assume that $f(t, \cdot)$ is not constant for any sufficiently small t, including $t = 0$. Then the dimension of S is one, and S is composed of finitely many analytic 1-simplexes, i.e. analytic paths given by $t = d_i(\tau)$, $w = c_i(\tau)$, where τ is the parameter in which c_i is analytic. If any of these simplexes do not pass through the $(0, p)$, decrease V to exclude them. The ones that do pass through the $(0, p)$ should be divided into two simplexes, each ending at $(0, p)$. Then we can assume that $\tau = 0$ corresponds to $(0, p)$, i.e. $d_i(0) = 0$ and $c_i(0) = p$. We have

$$f(d_i(\tau), c_i(\tau)) = 0.$$

Since d_i is analytic and not constant, it has only finitely many critical points. Decrease the size of V if necessary to exclude all nonzero critical points of d_i. Then either d_i is increasing in some interval $[0, b]$ or decreasing in some interval $[0, b]$. In either case it has an inverse function $\phi = d_i^{-1}$, so $d_i(\phi(t)) = t$. If the leading term of d_i is τ^n, then ϕ is real-analytic in $t^{1/n}$. We can parametrize the zero set of f by $\tilde{c}_i(t) = c_i(\phi(t))$, which is real-analytic in $t^{1/n}$. That completes the proof.

Remark: In applications it will usually be possible to replace the original parameter t by $t^{1/n}$, enabling us to assume that the zero set is analytic in t.

4 Eigenvalues of coverings of the sphere

In this section we present a result which will be applied to the Gauss map of a one-parameter family of minimal surfaces. However, we present it here as a result about conformal mappings from a plane domain to the Riemann sphere. We take the "Riemann sphere" to be the unit sphere $\{(x, y, z) : x^2+y^2+z^2 = 1\}$. Stereographic projection **St** is defined by

$$\mathbf{St}((x_1, x_2, x_3)) = \frac{x_1 + ix_2}{1 - x_3}$$

and its inverse is given by

$$\mathbf{St}^{-1}(z) = \frac{1}{1 + |z|^2} \begin{bmatrix} 2\,\mathrm{Re}\; z \\ 2\,\mathrm{Im}\; z \\ |z|^2 - 1 \end{bmatrix}.$$

Thus the equator projects onto the unit circle, and the "north pole" $(0, 0, 1)$ projects onto ∞, while the southern hemisphere projects onto the unit disk, with the south pole going to origin. The picture then has the plane passing through the equator of the sphere. We follow [23], p. 46, in these details, and we mention them because some other authors use a sphere of radius $1/2$, with the picture having the sphere entirely above the plane, tangent to the plane where the south pole of the sphere touches the origin of the plane.

Let D^+ be the upper half of the unit disk. We will consider a one-parameter family of analytic mappings from D^+ to the Riemann sphere. (These arise in our work as the Gauss maps of a one-parameter family of minimal surfaces, but here we consider a more abstract situation.) We write t for the parameter and $N(t, z)$ for the value of the mapping. For brevity we often omit the explicit t-dependence and write $N(z)$, or $N^t(z)$. Thus N^t is the map $N(t, \cdot)$ from D^+ to the Riemann sphere, for a fixed t. We define $g = \mathbf{St} \circ N$. The map N is thus conformal except at the zeros of g' and poles of g of order more than one. (These are sometimes known as *ramification points* of g.)

We will suppose that for sufficiently small t, the map g is a quotient of functions B/A, where B and A are analytic jointly in t and z (in the sense explained in the previous section), and that when $t = 0$, A has a zero of order m, and B has a zero of order $m + k$, for some positive integers m and k. (In our work, these arise from a boundary branch point of order $2m$ and index k, but here it is not necessary to be so specific.)

By Lemma 1, near the origin there exist paths $a_i(t)$ and $b_i(t)$ describing the poles and zeroes of g, respectively, so that each a_i and b_i is analytic in a rational power of t. Let s be the greatest common divisor of the denominators of these rational powers, and replace t by $t^{1/s}$. Then a_i and b_i will be analytic in t. We suppose that the parameter t has been chosen such that a_i and b_i are analytic in t. We also may suppose without loss of generality that for all sufficiently small positive t, we have $a_i(t) \neq b_j(t)$ for all i, j, since otherwise by analyticity we would have, for some i and j, $a_i(t) = b_j(t)$ for all sufficiently small t, and then the factor $z - a_i(t)$ could be cancelled out of both A and B. The partial derivatives of a quotient of real analytic functions are again quotients of real-analytic functions, so the structure of the set of critical points of the real and imaginary parts of such a function is also known: it is the union of a finite set of real-analytic arcs.

Let $A(t)$ be the area of the spherical image of D^+ under N, counting multiple coverings. Specifically, we have

$$\begin{aligned} A(t) &= \int\int_{D^+} \frac{1}{2}|\nabla N|^2 \, dx \, dy \\ &= \int\int_{D^+} \left[\frac{4|g'|}{(1+|g|^2)^2}\right]^2 dx \, dy \end{aligned}$$

Associated with a map N defined on a region Ω in the plane, there is a natural eigenvalue problem:[3]

$$\begin{aligned} \triangle\phi + \frac{1}{2}\lambda|\nabla N|^2\phi &= 0 \qquad \text{in } \Omega \\ \phi &= 0 \qquad \text{on } \partial\Omega \end{aligned}$$

We denote the least eigenvalue λ of this problem by $\lambda_{N,\Omega}$, or simply by λ_Ω when N is clear from the context. (In this work, we have no occasion to consider eigenvalues other than the least one.) The factor (1/2) arises because the area element on the sphere is $(1/2)|\nabla N|^2 dx \, dy$. The least eigenvalue is well-known to be equal to the infimum of the Rayleigh quotient

$$R[\phi] = \frac{\int\int_\Omega |\nabla\phi|^2 \, dx \, dy}{\int\int_\Omega (1/2)|\nabla N|^2\phi^2 \, dx \, dy}.$$

When we speak of the least eigenvalue λ_Ω of a region Ω on the Riemann sphere, we mean the following: Let Δ be the stereographic projection of Ω and N the inverse of stereographic projection. Then $\lambda_\Omega := \lambda_{N,\Delta}$. The eigenvalue problem $\triangle\phi - \frac{1}{2}\lambda|\nabla N|^2\phi$ on Δ is equivalent to the problem $\triangle\phi = \lambda\phi$ on Ω, where now $\triangle$ is the Laplace-Beltrami operator on the sphere. If Ω contains the north pole, we should use stereographic projection from some point not contained in Ω. We do not need to discuss the case when Ω is the entire sphere.

Example. We compute the least eigenvalue when $N(\Omega)$ is a hemisphere. In this case the eigenfunction in the lower hemisphere is minus the Z-component of N. For example with Ω equal to the unit disk and $g(z) = z$, we have $N(z)$ the inverse of stereographic projection. With $|z| = r$ and $z = x + iy$ we have

$$N(z) = \frac{1}{1+r^2}\begin{bmatrix} 2x \\ 2y \\ r^2 - 1 \end{bmatrix}.$$

The eigenfunction ϕ is given, with $|z| = r$, by

$$\phi(z) = \frac{1-r^2}{1+r^2}.$$

---

[3] As is discussed in [2] and [22], p. 103, among many other sources, when N is the Gauss map of a minimal surface, this eigenvalue problem is connected to the second variation of area. This does not concern us directly in this section. Some readers may be more familiar with another form of the eigenvalue equation for which the critical eigenvalue is zero rather than 2; both forms are discussed in [22], p. 103, cf. equations (62) and (62′).

A few lines of elementary computations (or a couple of commands to a computer algebra program) show that

$$\begin{aligned}\triangle\phi &= \phi_{rr} + \frac{1}{r}\phi_r \\ &= \frac{8(r^2-1)}{(1+r^2)^3}\end{aligned}$$

and

$$\begin{aligned}\frac{1}{2}|\nabla N|^2 &= \frac{1}{2}(N_x^2+N_y^2) = \frac{4r}{(1+r^2)^2} \\ \frac{1}{2}|\nabla N|^2\phi &= \frac{4(1-r^2)}{(1+r^2)^3}\end{aligned}$$

Hence, $\triangle\phi + |\nabla N|^2\phi = 0$, which means the eigenvalue of a hemisphere is 2.

Lemma 2 *Let Ω be a connected open set on the sphere with least eigenvalue λ_Ω. Suppose Δ is a region (open set) in the plane and $\Omega \subseteq N(\Delta)$ and $N(\partial\Delta)\cap\Omega = \phi$. Suppose that boundary of Ω is C^2 and the boundary of Δ is piecewise C^2 with the pieces meeting at positive angles. Then $\lambda_{N,\Delta} \le \lambda_\Omega$, and strict inequality holds if $N^{-1}\partial\Omega$ contains an interior point of Δ.*

Remark. Regarding the assumptions on the boundaries, the proof requires that the least eigenvalue be the minimum of the Rayleigh quotient, and that the gradient of the least eigenfunction of Ω not vanish at any boundary point. The hypotheses given imply these conditions but still allow Δ to be a half-disk. See [13].

Proof. Let ϕ be the least eigenfunction of Ω and define ψ on Δ by setting $\psi(z) = \phi(N(z))$ if $N(z) \in \Omega$ else $\psi(z) = 0$. Then ψ is admissible in the Rayleigh quotient for Δ, since ϕ is zero outside Ω and $N(\partial\Delta)\cap\Omega = \phi$. The Rayleigh quotient in question is

$$\frac{\int\int_\Delta |\nabla\psi|^2 dx\, dy}{\int\int_\Delta \frac{1}{2}|\nabla N|^2\psi^2}$$

(The factor 1/2 was explained above.) Since $\Omega \subseteq N(\Delta)$, on the support of ψ, N is a covering map (i.e., locally a homeomorphism), except at the points of ramification of N, which are isolated. Since Ω is connected and $N(\partial\Delta)\cap\Omega = \phi$, the number of sheets over (cardinality of the pre-image of) N of each non-ramification point is the same. Hence each of the two integrals in the (numerator and denominator of the) Rayleigh quotient is the number of sheets times the corresponding integral on the Riemann sphere, with ϕ in place of ψ. That is, the Rayleigh quotient for ψ on Δ equals the Rayleigh quotient for ϕ on Ω, which is λ_Ω. Since λ_Δ is the minimum of such Rayleigh quotients, $\lambda_\Delta \le \lambda_\Omega$. Now suppose there is an interior point p of Δ in $N^{-1}(\partial\Omega)$. By Hopf's lemma, $\nabla\phi$ is never zero at a point on $\partial\Omega$ where the boundary is C^2. By analyticity, ∇N is zero only at isolated points; so there is a point q near p which is still on $N^{-1}(\partial\Omega)$ at which ∇N is not zero and $\nabla\psi$ is not zero. Hence near q, the set $N^{-1}(\partial\Omega)$ is a smooth arc, and $\nabla\psi$ is zero on one side of it and bounded away from zero on the other side. Hence we can "smooth out the edge" near q to obtain a function ψ' which is admissible for the Rayleigh quotient and has smaller Rayleigh quotient than ψ. Hence $\lambda_\Delta < \lambda_\Omega$. This completes the proof of the lemma.

Remark. The lemma shows that if the Gaussian image of a minimal surface defined in Δ contains a hemisphere, then the eigenvalue $\lambda_{N,\Delta}$ is less than 2.

5 Basic Setup and Weierstrass Representation

We suppose we are in the following situation: a real-analytic Jordan curve, not lying in a plane, bounds a one-parameter family of minimal surfaces $u(t,z)$. The surfaces are parametrized by z in the upper half plane, so that on the real line each $u(t,\cdot)$ is a reparametrization of Γ. The surface $u(0,\cdot)$ has a boundary branch point of order $2m$ and index k at the origin. The surfaces $u(t,\cdot)$ for $t>0$ are immersed. All our arguments will be local; we shall only be concerned with what happens near the origin.[4]

We suppose that Γ passes through the origin tangent to the X-axis. Then since Γ is a Jordan curve, it is not contained in a line. We still are free to orient the Y and Z axes. We do this in such a way that the normal to $u(0,\cdot)$ at the branch point (which is well-defined) points in the positive Z-direction.

Finally, we can assume that for all t we have $u(0,0)=0$. If this is not already the case, then we can make it so by applying (for each $t>0$) a conformal transformation that moves $u_t(0)$ to the origin. If we do this while using the upper half-plane for a parameter domain, then the transformation in question is simply the map $z \mapsto z - \xi(t)$, where $u(\xi(t)) = 0$. $\xi(t)$ can be found analytically in a rational power of t, and we can replace t by a new parameter in which the family will be analytic. This assumption will be used Ifor example) to know that u can be found on the boundary by integrating u_x starting from 0.

The surfaces $u(t,z)$ have an Enneper-Weierstrass representation

$$u(t,z) = \mathrm{Re}\begin{bmatrix} \frac{1}{2}\int f - fg^2\,dz \\ \frac{i}{2}\int f + fg^2\,dz \\ \int fg\,dz \end{bmatrix}$$

The branch points of $u(t,\cdot)$ are the places where f and fg^2 vanish simultaneously. In [2] we were concerned with interior branch points only. In that case we can argue that since there are no branch points for $t>0$, the zeroes of f are of even order and coincide with the poles of g, so that f and g have the forms stated in Lemma 3.3 of [2], namely:

---

[4] We do not need to assume that the branch point is a true branch point. We can rule out the existence of a one-parameter family of the type considered without that assumption. This does not, however, provide a new proof of the non-existence of true branch points which are not the terminus of a one-parameter family of relative minima of area.

$$\begin{aligned} f(z) &= f_0(z)\prod_{i=1}^{m}(z-a_i)^2 \\ g(z) &= \frac{h(z)\displaystyle\prod_{i=1}^{m+k}(z-b_i)}{\displaystyle\prod_{i=1}^{m}(z-a_i)} \end{aligned}$$

However, this lemma was also used in [3], where boundary branch points are considered, and the formula was not established in this case. In this paper we correctly consider also the possibility that f and fg^2 may have some common zeroes $s_i(t)$, corresponding to a branch point (or branch points) in the lower half plane that converge to the origin as t goes to zero.

In general we might, *a priori*, have some common zeroes $s_i(t)$ of f and fg^2. These common zeroes of f and fg^2 could occur with different multiplicities in f and fg^2. Say that s_i occurs with multiplicity m_i in f and n_i in fg^2. We must have $\mathrm{Im}(s_i(t)) < 0$ for $t > 0$, since there will be branch points at the c_i, and $s_i(t)$, like $a_i(t)$ and $b_i(t)$, depends analytically on some power of t called t^γ and converging to 0 as t approaches 0.

We define

$$S(z) = \prod_{i=1}(z-s_i)^{\min(n_i,m_i)}$$

Then all zeroes of S are zeroes of f andfg^2 with at least the multiplicity they have in S. If s_i occurs with a greater multiplicity in fg^2 than in f, the extra occurrences are zeroes of g^2, and hence of g, so $n_i - m_i$ is even. Similarly, if s_i occurs with a greater multiplicity in f than in fg^2, the extra occurrences are poles of g^2, and hence poles of g, so $m_i - n_i$ is even. We have

$$\frac{fg^2}{f} = g^2$$

so the remaining zeroes of f and fg^2 are double. We can therefore define functions A and B, real-analytic in t and complex-analytic in z, such that

$$\begin{aligned} f(z) &= A^2S \\ f(z)g^2(z) &= -B^2S \\ g(z) &= \frac{fg^2}{f} = \frac{iB}{A} \\ f(z)g(z) &= iABS \end{aligned}$$

We already define the $a_i(t)$ and $b_i(t)$ to be the poles and zeroes of g, respectively. Let $2N$ be the number of zeros of S. (It must be even since the number of zeroes of f is $2m$ and all the zeroes of f besides those of S are double.) Then we can find functions A_0 and B_0, real-analytic in t and complex-analytic in z, not vanishing at the origin, such that

$$\begin{aligned} A(z) &= A_0(z,t)\prod_{i=1}^{m-N}(z-a_i) \\ B(z) &= i\lambda B_0(z,t)\prod_{i=1}^{m+k-N}(z-b_i) \end{aligned}$$

and $A_0(0,0) = 1$, $B_0(0,0)$ is real, and $|\lambda| = 1$ with $\lambda \neq -1$. Note that the common zeroes s_i do not enter into the formula for g. In [3], in effect we assumed the s_i, and hence S, do not occur. We can assume $A_0(0,0) = 1$, since we can always change z to a constant times z. The leading term of $B(z)$, namely $i\lambda B_0(0,0)$ is some nonzero complex constant, but because of the presence of λ in the formula, we can assume $B_0(0,0)$ is real.

Note that with this definition of A and B, some of the a_i and/or some of the b_i may have been defined to be equal to some of the s_i, in case s_i occurs with a greater multiplicity in f than in fg^2 or vice-versa. This is possible since these extra occurrences are double, since they occur as roots or poles of g. Each branch point s_i will either occur as an a_i, or as a b_i, or neither one (in case its multiplicities in f and fg^2 are equal); so it will still be true that $a_i(t)$ is not identically equal to $b_j(t)$ as t varies, even if the s_i occur and are treated this way.

As we have already remarked, it follows from Lemma 1 that the a_i, b_i, and s_i depend analytically on t. Although they all approach zero as t goes to zero, some may go to zero faster than others. The roots a_i, b_i, and s_i occur in “rings”, going to zero at different speeds. Ring 1 contains the principal roots. There is a (finite) sequence of powers $\gamma_1 < \gamma_2 < \ldots$ such that every root goes to zero as t^{γ_n} for some n; we say those roots belong to the n-th ring. Specifically, the roots in the n-th ring are those for which there exist nonzero complex constants α_i, β_i, and ζ_i such that

$$\begin{aligned} a_i &= \alpha_i t_n^\gamma + O(t^{\gamma+1}) \\ b_i &= \beta_i t_n^\gamma + O(t^{\gamma+1}) \\ s_i &= \zeta_i t_n^\gamma + O(t^{\gamma+1}) \end{aligned}$$

The roots in the first ring, i.e. those that go to zero as t^γ, are called the *principal roots.*

6 The Gauss map and the w-plane

Suppose that γ is one of the numbers γ_n; that is, γ is a number such that some of the roots a_i, b_i, or s_i go to zero as t^γ, but not necessarily the least such number; thus the principal roots go to zero more slowly that t^γ. It might be that $\gamma = \gamma_1$, in which case the roots that go to zero as t^γ are the principal roots;

but we are also interested in the case when $\gamma > \gamma_1$. We introduce notation as follows:

$$Q \text{ is the number of } a_i \text{ that go to zero as } t^\gamma \text{ or faster.} \tag{2}$$

$$S \text{ is the number of } s_i \text{ that go to zero as } t^\gamma \text{ or faster.} \tag{3}$$

$$R \text{ is the number of } b_i \text{ that go to zero as } t^\gamma \text{ or faster.} \tag{4}$$

Thus if $\gamma = \gamma_1$, we have $Q = m$, $S = 2N$, $R = m + k$, and $K = k$. Recall that $z = a_i(t)$ are the zeroes of A and $z = b_i(t)$ are the zeroes of B. We will generally be considering only one "ring" of roots at a time, and when we speak of α_i, β_i, and ζ_i, we have in mind those roots that go to zero as t^γ:

$$\begin{aligned} a_i &= \alpha_i t^\gamma (1 + O(t)) \\ b_i &= \beta_i t^\gamma (1 + O(t)) \\ a_i &= \zeta_i t^\gamma (1 + O(t)) \end{aligned}$$

We will refer to a_i, b_i, and s_i that go to zero as lower powers of t than t^γ as "slow" roots, and those that go to zero as higher powers of t as "fast" roots. We have

$$A = A_0(t, z) \prod_{i=1}^{m-N} (z - a_i(t))$$

We introduce a new variable w by $z = t^\gamma w$. Then a factor $z - \alpha_i t^\gamma$ becomes $t^\gamma(w - \alpha_i)$, and a factor $z - \alpha_i t^{\gamma_i}$ for $\gamma_i < \gamma$ becomes

$$t^{\gamma_i}(z t^{\gamma - \gamma_i} - \alpha_i) = t^{\gamma_i}(-\alpha_i + O(t)).$$

Thus the slow roots introduce constants and powers of t, and the roots that go to zero as t^γ introduce factors containing $w - \alpha_i$, and the roots that go to zero faster than t^γ introduce factors $w(1 + O(t))$. All products without explicit indices will be understood to be over those roots which are $O(t^\gamma)$, and in such products (but not elsewhere) we will assume that the α_i corresponding to the fast roots are zero. Thus $\prod(w - \alpha_i)$ contains a factor of w for each fast root, and the total number of factors is the number of a_i that go to zero as t^γ or faster. Define, for the roots that go to zero as t^γ or faster,

$$\begin{aligned} \alpha &:= \text{the product of all the } -\alpha_i \text{ over the slow } a_i \\ \beta &:= \text{the product of the } -\beta_i \text{ over the slow } b_i \\ \zeta &:= \text{the product of the } -\zeta_i \text{ over the slow } \zeta_i \\ \mathbf{a}_i &:= a_i / t^\gamma \\ \mathbf{b}_i &:= b_i / t^\gamma \\ \mathbf{s}_i &:= s_i / t^\gamma \\ \tilde{\mathbb{A}} &:= \alpha \prod (w - \mathbf{a}_i) \qquad \text{(the product is over } i \text{ such that } a_i = O(t^\gamma)) \\ \tilde{\mathbb{B}} &:= B_0 \beta \prod (w - \mathbf{b}_i) \\ \tilde{\mathbb{S}} &:= \zeta \prod (w - \mathbf{s}_i) \end{aligned}$$

The tilde indicates t-dependence. When $t = 0$ we drop the tilde:

$$\begin{aligned} \mathbb{A} &= \alpha \mathbb{A}_0(t,w) \prod (w - \alpha_i) \\ \mathbb{B} &= B_0 \beta \mathbb{B}_0(t,w) \prod (w - \beta_i) \\ \mathbb{S} &= \zeta \prod (w - \zeta_i) \end{aligned}$$

where $\mathbb{A}_0$ and $\mathbb{B}_0$ are real analytic in t and complex analytic in w, zero at the origin, arising from A_0 and B_0 which are real analytic in t and complex analytic in z.

$$A = t^d t^{Q\gamma} \tilde{\mathbb{A}}(1 + O(t)) \tag{5}$$

The power d is the sum of the γ_i, one γ_i for each a_i that goes to zero as γ_i for $\gamma_i > \gamma$. Similarly, we have numbers e and f such that

$$\begin{aligned} B &= t^e t^{R\gamma} \tilde{\mathbb{B}}(1 + O(t)) \qquad (6) \\ S &= t^f t^{S\gamma} \tilde{\mathbb{S}}(1 + O(t)) \qquad (7) \end{aligned}$$

No ambiguity will result from using f for this power of t since in this meaning, it will always occur in an exponent, and the Weierstrass function f never occurs in an exponent. We define

$$\mathcal{K} := (R - Q)\gamma + (e - d). \tag{8}$$

When $\gamma = \gamma_1$, so we are working with the principal, or first-ring, roots, then $\mathcal{K} = k\gamma_1$.

We review the formula for the unit normal. The function g in the Weierstrass representation, which is the stereographic projection of the unit normal N, is given by $g(z) = iB/A$, and the normal itself is given by

$$N = \frac{1}{|g|^2 + 1} \begin{bmatrix} 2\mathrm{Re}\,(g) \\ 2\,\mathrm{Im}(g) \\ |g|^2 - 1 \end{bmatrix}. \tag{9}$$

We write $\bar{A}$ for the complex conjugate of A. Substituting $g = iB/A$, we have

$$N = \frac{1}{1 + |B/A|^2} \begin{bmatrix} -2\mathrm{Im}\,(B/A) \\ 2\mathrm{Re}(B/A) \\ -1 + |B/A|^2 \end{bmatrix}$$

Expressing this in terms of $\tilde{\mathbb{B}}/\tilde{\mathbb{A}}$, we have $\tilde{\mathbb{B}}/\tilde{\mathbb{A}} = t^{\mathcal{K}} B/A$, and hence

$$N = \frac{1}{1 + t^{2\mathcal{K}} |\tilde{\mathbb{B}}/\tilde{\mathbb{A}}|^2} \begin{bmatrix} -2\mathrm{Im}\,(\tilde{\mathbb{B}}/\tilde{\mathbb{A}}) \\ 2\mathrm{Re}\,(\tilde{\mathbb{B}}/\tilde{\mathbb{A}}) \\ -1 + t^{2\mathcal{K}} |\mathbb{B}/\mathbb{A}|^2 \end{bmatrix} \tag{10}$$

It follows that

$$\frac{{}^1N}{{}^2N} = -\frac{\mathrm{Im}\,(\tilde{\mathbb{B}}/\tilde{\mathbb{A}})}{\mathrm{Re}\,(\tilde{\mathbb{B}}/\tilde{\mathbb{A}})} \tag{11}$$

Lemma 3 *Suppose that on compact subsets of the w-plane away from the zeroes of $\mathbb{A}$, N converges to $(-1,0,0)$ as t goes to zero. Then*

(i) $\mathbb{A}$, $\mathbb{S}$, and $\mathbb{B}$ are real;

(ii) the α_i and ζ_i are real and the β_i are real and/or occur in complex-conjugate pairs.

Remark. The hypothesis is certainly true for the first ring of a_i, the principal roots. We will later show inductively that it is true for all the rings of a_i, i.e. when each γ_n is used to define w.

Proof. First we prove that α_i does not have positive imaginary part. Suppose, for proof by contradiction, that it does. Then consider a circle of radius ρ about α_i, where ρ small enough that the entire circle lies in the upper half plane and each other α_i is either inside or outside the circle. Then by hypothesis, on the boundary of the circle, N goes to $(-1,0,0)$ as t goes to zero. Hence the Gauss map covers the upper hemisphere inside the circle, so the eigenvalue is less than 2, contradiction. That contradiction proves that α_i does not have positive imaginary part.

Let $Y(w) := {}^2u(t^\gamma w)$. Then

$$Y(w) = \frac{C_2}{q+1}\tau^{q+1}(1+O(s))$$

where τ is as in (1).

$$\begin{aligned}\tau &= X(w) = \int_0^z A^2S + B^2S\,dz \\ &= t^{2d+2Q\gamma+f+S\gamma+1}\int_0^w \mathbb{A}^2\mathbb{S} + t^{2\mathcal{K}\gamma}\mathbb{B}^2\,dw(1+O(t))\end{aligned}$$

by (5),(7), (6), and (8). The condition that N converges to $(-1,0,0)$ away from the zeroes of $\mathbb{A}$ can be succinctly expressed as $\mathcal{K} > 0$, since the stereographic projection of the normal is given by $t^{\mathcal{K}}\tilde{\mathbb{B}}/\tilde{\mathbb{A}}$. Hence the term in $t^{\mathcal{K}}$ can be absorbed into the error term, and we have

$$\tau = t^{2d+2Q\gamma+f+S\gamma+1}\int_0^w \mathbb{A}^2\mathbb{S}\,dw(1+O(t))$$

and hence

$$Y(w) = \frac{C_2}{q+1}t^{(q+1)(2d+2Q\gamma+f+S\gamma+1)}\int_0^w \mathbb{A}^2\mathbb{S}\,dw(1+O(t))$$

On the other hand

$$Y(w) = t^{(2d+2Q\gamma+f+S\gamma+1)}\,\mathrm{Im}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw + O(t^{(2d+2Q\gamma+f+S\gamma+2)})$$

It follows that

$$\mathrm{Im}\int_0^w \mathbb{A}^2\mathbb{S}\,dw = 0$$

since it is independent of t but from the two previous equations must be $O(t)$, since $q > 0$. Therefore the polynomial $\int_0^w \mathbb{A}^2\mathbb{S}\,dw$ is real on the real axis. It follows that its coefficients are real. Hence the coefficients of its derivative $\mathbb{A}^2\mathbb{S}$ are real. Hence the roots of $\mathbb{A}^2\mathbb{S}$, which are the α_i and ζ_i, come in complex-conjugate pairs. But we proved above that no α_i has positive imaginary part. Hence they are all real when $t = 0$. Similarly, the ζ_i do not have positive imaginary part, since u^t has no branch points for $t > 0$. Hence the ζ_i are real when $t = 0$.

Similarly, with $Z(w) = {}^3u(t^\gamma)w$ in place of Y, we find

$$\operatorname{Im}\int_0^w \mathbb{A}\mathbb{B}\mathbb{S}\,dw = 0$$

That is, $\int_0^w \mathbb{A}\mathbb{B}\mathbb{S}\,dw = 0$ is real on the real axis. Therefore also its derivative $\mathbb{A}\mathbb{B}\mathbb{S}$ is real on the real axis. But $\mathbb{A}\mathbb{B}\mathbb{S}$ is a polynomial; since it is real on the real axis, its coefficients are real. We have already proved that $\mathbb{A}\mathbb{S}$ is real on the real axis; hence $\mathbb{B}$ is also real on the real axis. Hence its roots, which are the $\hat{\mathbf{b}}_i$ when $t = 0$, are real or occur in complex-conjugate pairs. That completes the proof of the lemma.

7 The eigenfunction

In this section, we compute the eigenfunction of the variational problem associated with the second variation of area. This is shown in [2] to be $\phi = u_t \cdot N$, where $u_t = du/dt$ is the "tangent vector" to the one-parameter family, and N is the unit normal. The calculations in this section are similar to those in [3], but applying to the whole w-plane rather than only to compact subsets away from the α_i, and applying not in the vicinity of the roots going to zero at the slowest rate, but also to the inner "rings" of roots.

Only two assumptions will be made for our eigenfunction computation: we assume that $e > d$ and $R > Q$. (See (5), (6), (7), and (8) for the definitions of these quantities.) These assumptions imply that the power of t that goes with $\mathbb{B}$ is larger than the power of t that goes with $\mathbb{A}$. In the case $\gamma = \gamma_1$, we have $b = d = 0$ and $R = m + k - N$ and $Q = m - N$, so the inequality in question is $m + k - N > m - N$. This assumption ensures that the unit normal, whose stereographic projection is B/A, is equal to t to a positive power time $\mathbb{B}/\mathbb{A}$, so that on compact subsets of the w-plane away from zeroes of $\mathbb{A}$, the unit normal tends to $(-1, 0, 0)$. Specifically we have

$$\begin{aligned}\frac{B}{A} &= t^{(R-Q)\gamma+(e-d)}\frac{\mathbb{B}}{\mathbb{A}} \\ &= t^{\mathcal{K}\gamma}\frac{\mathbb{B}}{\mathbb{A}} \qquad \text{by (8)}\end{aligned}$$

and the exponent of t is positive, by our assumptions. Differentiating the Weierstrass representation for u, we have

$$2u_t = \frac{d}{dt}\mathrm{Re}\begin{bmatrix} \int_0^z (A^2S + B^2S)\,dz \\ i\int_0^z (A^2S - B^2S)\,dz \\ 2i\int_0^z ABS\,dz \end{bmatrix}$$

Expressing this in terms of $\tilde{\mathbb{A}}$ and $\tilde{\mathbb{B}}$ we have

$$2u_t = \frac{d}{dt}\mathrm{Re}\begin{bmatrix} t^{2d+f}t^{(2Q+S+1)\gamma}\int_0^w (\tilde{\mathbb{A}}^2\tilde{\mathbb{S}} + t^{2\mathcal{K}}\tilde{\mathbb{B}}^2\tilde{\mathbb{S}})\,dw(1+O(t)) \\ it^{2d+f}t^{(2Q+S+1)\gamma}\int_0^w (\tilde{\mathbb{A}}^2\tilde{\mathbb{S}} - t^{2\mathcal{K}}\tilde{\mathbb{B}}^2\tilde{\mathbb{S}})\,dw(1+O(t)) \\ 2it^{2d+f}t^{(2Q+S+1)\gamma+\mathcal{K}}\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,dw(1+O(t)) \end{bmatrix}$$

where the power of t in the last term is calculated thus:

$$e+d+f+(R+Q+S+1)\gamma \quad = \quad 2d+f+(2Q+S+1)\gamma+\mathcal{K} \qquad \text{by (8)}$$

Since the expressions for these exponents are a bit unwieldy, we shorten them by introducing

$$\mathcal{M} \quad := \quad 2d+f+(2Q+S+1)\gamma \tag{12}$$

Then we have

$$\begin{align} \int_0^w \tilde{\mathbb{A}}^2\,dw &= t^{\mathcal{M}}\int_0^z A^2\,dz \tag{13} \\ \int_0^w \tilde{\mathbb{B}}^2\,dw &= t^{\mathcal{M}+\mathcal{K}}\int_0^z B^2\,dz \tag{14} \\ & \tag{15} \end{align}$$

The equation for u_t becomes

$$2u_t = \frac{d}{dt}\mathrm{Re}\begin{bmatrix} t^{\mathcal{M}}\int_0^w (\tilde{\mathbb{A}}^2\tilde{\mathbb{S}} + t^{2\mathcal{K}}\tilde{\mathbb{B}}^2\tilde{\mathbb{S}})\,dw(1+O(t)) \\ it^{\mathcal{M}}\int_0^w (\tilde{\mathbb{A}}^2\tilde{\mathbb{S}} - t^{2\mathcal{K}}\tilde{\mathbb{B}}^2\tilde{\mathbb{S}})\,dw(1+O(t)) \\ 2it^{\mathcal{M}+\mathcal{K}}\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,dw(1+O(t)) \end{bmatrix}$$

The rule for differentiating with respect to t in this situation is found by applying the chain rule as follows, where d/dt means the partial derivative with respect to t, holding z fixed, and $\partial/\partial t$ means the partial derivative with respect to t, holding w fixed.

$$\begin{aligned} dH/dt &= \frac{\partial H}{\partial t} + \frac{dH}{dw}\frac{dw}{dt} \\ &= \frac{\partial H}{\partial t} + \frac{dt^{-\gamma}z}{dt}\frac{dH}{dw} \\ &= \frac{\partial H}{\partial t} - \gamma t^{-\gamma-1}z\frac{dH}{dw} \\ &= \frac{\partial H}{\partial t} - \gamma t^{-\gamma-1}t^{\gamma}w\frac{dH}{dw} \end{aligned}$$

The final result for differentiating with respect to t is then

$$\frac{dH}{dt} = \frac{\partial H}{\partial t} - t^{-1}\gamma w \frac{dH}{dw} \tag{16}$$

Applying this rule we find, omitting for legibility the $(1 + O(t))$ factors that should multiply every term,

$$\begin{aligned}
&2u_t = \\
&\operatorname{Re}\begin{bmatrix} t^{\mathcal{M}-1}\mathcal{M}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw + t^{\mathcal{M}+2\mathcal{K}-1}(\mathcal{M}+2\mathcal{K})\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,dw \\ it^{\mathcal{M}-1}\mathcal{M}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw + it^{\mathcal{M}+2\mathcal{K}-1}(\mathcal{M}+2\mathcal{K})\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,dw \\ 2it^{\mathcal{M}+\mathcal{K}-1}(\mathcal{M}+\mathcal{K})\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,dw \end{bmatrix} \\
&-\operatorname{Re}\begin{bmatrix} t^{\mathcal{M}-1}\gamma\tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,w + t^{\mathcal{M}+2\mathcal{K}-1}\tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,w \\ it^{\mathcal{M}-1}\gamma\tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,w + it^{\mathcal{M}+2\mathcal{K}-1}\tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,w \\ 2it^{\mathcal{M}+\mathcal{K}-1}\gamma\tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,w \end{bmatrix} \\
&+\operatorname{Re}\begin{bmatrix} t^{\mathcal{M}}\mathcal{M}\frac{\partial}{\partial t}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw + t^{\mathcal{M}+2\mathcal{K}}(\mathcal{M}+2\mathcal{K})\frac{\partial}{\partial t}\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,w \\ it^{\mathcal{M}}\mathcal{M}\frac{\partial}{\partial t}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw + it^{\mathcal{M}+2\mathcal{K}}(\mathcal{M}+2\mathcal{K})\frac{\partial}{\partial t}\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,dw \\ 2it^{\mathcal{M}+\mathcal{K}}(\mathcal{M}+\mathcal{K})\frac{\partial}{\partial t}\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,dw \end{bmatrix}
\end{aligned}$$

Note that the last terms, involving $\partial/\partial t$, have one higher power of t and are in some sense “error terms” that were legitimately neglected when considering convergence on compact subsets away from the α_i. In addition to those terms there is also the term $O(1)u_t$ resulting from differentiating the $(1+O(t))$ factor. That term has one more power of t than the terms shown, and can be absorbed into the $(1 + O(t))$ factors multiplying each term.

Combining the first two vectors into one, and dividing by the leading power of t, the unwritten factors of $(1 + O(t))$ and the terms involving $\partial/\partial t$ can be expressed as the $O(t)$ in the following formula:

$$\begin{aligned}
&\frac{2u_t}{t^{\mathcal{M}-1}} = \\
&\operatorname{Re}\begin{bmatrix} \mathcal{M}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw - \gamma\tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,w + t^{2\mathcal{K}}((\mathcal{M}+2\mathcal{K})\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,dw - \gamma\tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,w) + O(t) \\ (\mathcal{M}i\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw - i\gamma\tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,w + it^{2\mathcal{K}}((\mathcal{M}+2\mathcal{K})\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,dw - \gamma\tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,w) + O(t) \\ 2it^{\mathcal{K}}(\mathcal{M}+\mathcal{K})\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,dw - \gamma\tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,w) + O(t) \end{bmatrix}
\end{aligned}$$

Taking the dot product with N, we have the following formula for the eigenfunction, valid in the whole w-plane, including subsets that contain the zeroes of $\mathbb{A}$:

$$\begin{aligned}
&\frac{2}{t^{\mathcal{M}-1}}\phi = \\
&[{}^1N]\operatorname{Re}\left(\mathcal{M}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw - \gamma\tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,w + O(t)\right) \\
&+t^{2\mathcal{K}}[{}^1N]\operatorname{Re}\left((\mathcal{M}+2\mathcal{K})\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,dw - \gamma\tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,w + O(t)\right)
\end{aligned} \tag{17}$$

$$\begin{aligned}
&-[{}^{2}N]\,\mathrm{Im}\Bigg(\mathcal{M}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,w+O(t)\Bigg)\\
&-t^{2k\gamma}[{}^{2}N]\,\mathrm{Im}\Bigg((\mathcal{M}+2\mathcal{K})\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,w+O(t)\Bigg)\\
&-t^{k\gamma}(\mathcal{M}+\mathcal{K})[{}^{3}N]2\,\mathrm{Im}\Bigg(\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,w+O(t)\Bigg)
\end{aligned}$$

We can now substitute for ${}^{1}N$, ${}^{2}N$, and ${}^{3}N$ to express the formula entirely in terms of $\tilde{\mathbb{B}}$ and $\tilde{\mathbb{A}}$. We get (after multiplying by the denominator and changing the signs of both sides)

$$\begin{aligned}
&\frac{-1(1+t^{2\mathcal{K}}|\tilde{\mathbb{B}}/\tilde{\mathbb{A}}|^2)}{t^{\mathcal{M}-1}}\phi=\\
&[t^{\mathcal{K}\gamma}\mathrm{Im}\,(\tilde{\mathbb{B}}/\tilde{\mathbb{A}})]\,\mathrm{Re}\Bigg(\mathcal{M}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,w+O(t)\Bigg)\\
&+t^{2\mathcal{K}\gamma}[t^{\mathcal{K}\gamma}\mathrm{Im}\,(\tilde{\mathbb{B}}/\tilde{\mathbb{A}})]\,\mathrm{Re}\Bigg((\mathcal{M}+2\mathcal{K})\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,w+O(t)\Bigg)\\
&+[t^{\mathcal{K}\gamma}\mathrm{Re}\,(\tilde{\mathbb{B}}/\tilde{\mathbb{A}})]\,\mathrm{Im}\Bigg(\mathcal{M}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,w+O(t)\Bigg)\\
&+t^{2\mathcal{K}\gamma}[t^{\mathcal{K}\gamma}\mathrm{Re}\,(\tilde{\mathbb{B}}/\tilde{\mathbb{A}})]\,\mathrm{Im}\Bigg(\mathcal{M}+2\mathcal{K})\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,w+O(t)\Bigg)\\
&+t^{\mathcal{K}\gamma}(\mathcal{M}+2\mathcal{K})\Bigg(-1+t^{2\mathcal{K}\gamma}|\tilde{\mathbb{B}}/\tilde{\mathbb{A}}|^2\Bigg)\Bigg(\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,w+O(t)\Bigg)
\end{aligned}$$

Combining real and imaginary parts and dividing by another $t^{\mathcal{K}}$ we find the formula in the following lemma.

Lemma 4 *The following formula for the least eigenfunction of u^t in terms of N is valid everywhere, even near the zeroes of $\mathbb{A}$:*

$$\begin{aligned}
&\frac{-(1+t^{2\mathcal{K}\gamma}|\tilde{\mathbb{B}}/\tilde{\mathbb{A}}|^2)}{t^{(\mathcal{M}+\mathcal{K})\gamma-1}}\phi=\\
&\mathrm{Im}\Bigg[(\tilde{\mathbb{B}}/\tilde{\mathbb{A}})\Bigg(\mathcal{M}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,w+O(t)\Bigg)\Bigg]\\
&-\Bigg(1-t^{2\mathcal{K}\gamma}|\tilde{\mathbb{B}}/\tilde{\mathbb{A}}|^2\Bigg)\mathrm{Im}\Bigg[(\mathcal{M}+\mathcal{K})\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,w+O(t)\Bigg]\\
&+t^{\mathcal{K}\gamma}\mathrm{Im}\Bigg[(\tilde{\mathbb{B}}/\tilde{\mathbb{A}})\Bigg((\mathcal{M}+2\mathcal{K})\int_0^w \tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,dw-\gamma\tilde{\mathbb{B}}^2\tilde{\mathbb{S}}\,w+O(t)\Bigg)\Bigg]
\end{aligned}$$

Proof. This was proved above.

Lemma 5 *The following formulas for the least eigenfunction of u^t in terms of N is valid for convergence on compact subsets of the w plane away from the* *zeroes of* $\mathbb{A}$*, provided* $t^{\mathcal{K}}\tilde{\mathbb{B}}/\tilde{\mathbb{A}} = O(t)$ *on those subsets.*

$$\begin{aligned}
&\frac{-1}{t^{(\mathcal{M}+\mathcal{K})\gamma-1}}\phi = \\
&\operatorname{Im}\left\{\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\left(\mathcal{M}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw - \gamma\tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,w\right) - (\mathcal{M}+\mathcal{K})\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,dw - \gamma\tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,w + O(t)\right\} \\
= \quad &\operatorname{Im}\left\{\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\left(\mathcal{M}\int_0^w \tilde{\mathbb{A}}^2\tilde{\mathbb{S}}\,dw - \right) - (\mathcal{M}+\mathcal{K})\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\tilde{\mathbb{S}}\,dw + O(t)\right\}
\end{aligned}$$

Remark. The provision that $t^{\mathcal{K}}\tilde{\mathbb{B}}/\tilde{\mathbb{A}} = O(t)$ is the same as providing that N approach $(-1, 0, 0)$ on those subsets. It is certainly true when $\gamma = \gamma_1$, and later we will inductively show it is true on the inner rings of roots as well.

Proof. The first formula of the lemma follows from the previous lemma by replacing $t^{\mathcal{K}}\tilde{\mathbb{B}}/\tilde{\mathbb{A}}$ with $O(t)$. The second formula follows by canceling two terms in the first formula.

We now define the function $\mathbb{H}$ by

$$\mathbb{H}(w) \quad = \quad \frac{\mathbb{B}}{\mathbb{A}}\mathcal{M}\int_0^w \mathbb{A}^2\mathbb{S}\,dw - (\mathcal{M}+\mathcal{K})\int_0^w \mathbb{A}\mathbb{B}\mathbb{S}\,dw \tag{18}$$

Then by the previous lemma we have

$$\operatorname{Im}\mathbb{H} \quad = \quad \lim_{t\to 0}\frac{-2}{t^{(\mathcal{M}+\mathcal{K})\gamma-1}}\phi$$

on compact subsets of the w-plane away from the zeroes of $\mathbb{A}$. A crucial question is whether $\mathbb{H}$ is constant or not. If it is constant, then all the visible terms cancel out, and we can thus get no information from them. If it is not constant, we can get some information by the asymptotic behavior of $\mathbb{H}$ near the origin or near α_i.

8 What is true if H is not constant

In this section, we suppose that γ is one of the γ_n, that $z = t^\gamma w$, and that N goes to $(-1, 0, 0)$ on compact subsets away from the zeroes of $\mathbb{A}$. This condition we call the "N-condition". Since

$$\frac{B}{A} = t^{\mathcal{K}}\frac{\mathbb{B}}{\mathbb{A}}$$

the N-condition can be succinctly expressed as $\mathcal{K} > 0$. The α_i are the coefficients of the a_i in this ring of roots, i.e., $a_i = \alpha_i t^\gamma(1 + O(t))$, and we define $\alpha_i = 0$ for roots that go to zero faster than t^γ. Similarly for the β_i and ζ_i. We refer to the roots that go to zero as t^γ as "principal roots", and the roots that go to zero faster as "fast roots".

Lemma 6 *Suppose the N-condition holds. Then*

(i) There is at least one principal α_i or ζ_i, i.e., at least one a_i or s_i goes to zero as t^γ.

(ii) If $\mathbb{H}$ is not constant, then no a_i or s_i goes to zero faster than t^γ.

Proof. Let P be the total number of fast roots. Let Q, S, and R be the total number of a_i, s_i, and b_i going to zero faster than t^γ. Let $\hat{\alpha}$ be the product of the nonzero α_i (all of which are real) over the a_i that go to zero as t^γ. Let $\hat{\beta}$ be the product of the nonzero β_i. If the β_i are not real, they come in complex conjugate pairs, by Lemma 3, so the product $\hat{\beta}$ is real. Let $\hat{\zeta}$ be the product of the nonzero ζ_i, which is real since all the ζ_i are real. (Do not confuse $\hat{\alpha}$ with α, which is the product of the α_i over the slow roots, i.e. the ones going to zero slower than t^γ.)

We wish to analyze the asymptotic behavior of $\mathbb{H}$ near the origin. We will show that, if $\mathbb{H}$ is not constant, its leading term is a nonzero real constant times βw^{P+1}. Once we have done that, we can proceed as follows: Since $\mathbb{H}$ is not constant, its imaginary part, which is a multiple of the least eigenfunction, must have only one sign in the upper half plane, and that is possible only if $P = 0$. Therefore, there are no fast roots, which is conclusion (ii) of the lemma.

It remains to prove that $\mathbb{H}$ has the stated asymptotic form. We analyze the first formula of Lemma 5, rather than the second one. Letting t go to zero in that formula and using the definition of $\mathbb{H}$ we have

$$\begin{aligned} \mathbb{H} = {} & \frac{\mathbb{B}}{\mathbb{A}}\left(\mathcal{M}\int_0^w \mathbb{A}^2\mathbb{S}\,dw - \gamma\mathbb{A}^2\mathbb{S}\,w\right) \\ & -(\mathcal{M}+\mathcal{K})\int_0^w \mathbb{A}\mathbb{B}\mathbb{S}\,dw - \gamma\mathbb{A}\mathbb{B}\mathbb{S}\,w \end{aligned} \tag{19}$$

Consider the second term in that equation, namely

$$(\mathcal{M}+\mathcal{K})\int_0^w \mathbb{A}\mathbb{B}\mathbb{S}\,dw - \gamma\mathbb{A}\mathbb{B}\mathbb{S}\,w.$$

The total number of "fast roots", i.e. roots going to zero faster than t^γ, is $Q+R+S$. At least one of the a_i or b_i or s_i is a principal root, that is, it goes to zero as t^γ, not faster. There the total number P of fast roots among the a_i, b_i, and s_i is strictly less that $Q+R+S$. Therefore it is not the case that $\mathbb{A}\mathbb{B}\mathbb{S} = O(w^{(Q+R+S)/\gamma})$. Indeed, asymptotically for small w, $\mathbb{A}\mathbb{B}\mathbb{S}$ is a constant times w^P. Hence, asymptotically for small w, this term has the form cw^{P+1} for some nonzero real constant c. The value of c is

$$c = \alpha\beta\zeta B_0\hat{\alpha}\hat{\beta}\hat{\zeta}\left[\frac{\mathcal{M}+K}{P+1} - \gamma\right].$$

(The constants $\hat{\alpha}$, etc., are defined in the first paragraph of this proof.) To see that c is not zero, we recall the definitions of $\mathcal{M}$ and $\mathcal{K}$:

$$\begin{aligned} \mathcal{M} &= 2d + f + (2Q+S+1)\gamma \qquad \text{by (12)} \\ \mathcal{K} &= (R-Q)\gamma + (e-d) \qquad \text{by (8)} \end{aligned}$$

Adding, we find $\mathcal{M}+\mathcal{K} \geq (Q+R+S+1)\gamma$, and since $P < Q+R+S$, we have $\mathcal{M}+\mathcal{K} > (P+1)\gamma$. Hence $c \neq 0$ as claimed. Summarizing our result so far we have

$$(\mathcal{M}+\mathcal{K})\int_0^w \mathbb{ABS}\,dw - \gamma\mathbb{ABS}\,w \quad = \quad cw^{P+1}$$

Now consider the behavior of the first term in (19) near the origin, namely

$$\frac{\mathbb{B}}{\mathbb{A}}\bigg(\mathcal{M}\int_0^w \mathbb{A}^2\mathbb{S}\,dw - \gamma\mathbb{A}^2\mathbb{S}\,w\bigg). \tag{20}$$

Suppose, for purposes of contradiction, that there are no principal a_i or s_i, that is, all the principal roots are among the b_i. Then $\hat{\alpha} = \hat{\zeta} = 1$ (so those factors do not appear in c) and

$$\mathcal{M}\int_0^w \mathbb{A}^2\mathbb{S}\,dw = \frac{\mathcal{M}\alpha^2\zeta}{2Q+S+1}w^{2Q+S+1}$$

and

$$\gamma\mathbb{A}^2\mathbb{S}w = \gamma\alpha^2\zeta w^{2Q+S+1}$$

so the first term of (19) comes to

$$B_0\beta\alpha\zeta\hat{\beta}\bigg(\frac{\mathcal{M}}{2Q+S+1} - \gamma\bigg)w^{2Q+S+1} \tag{21}$$

In the outer ring, where $\mathcal{M} = \gamma(2m+1) = (2Q+S+1)\gamma$, this would make the entire term vanish, but in the inner rings, $\mathcal{M}$ can be larger. Subtracting the contributions of the two terms as given in (21) and (20), we have

$$\operatorname{Im}\mathbb{H} \quad = \quad cw^{P+1} - B_0\beta\alpha\zeta\hat{\beta}\bigg(\frac{\mathcal{M}}{2Q+S+1} - \gamma\bigg)w^{2Q+S+1} \tag{22}$$

Suppose, for proof by contradiction, that these two terms cancel. Then $2Q+S = P$ (i.e. no b_i is a fast root), and the coefficients must cancel too. That means

$$0 = \alpha\beta\zeta\hat{\beta}B_0\bigg[\frac{\mathcal{M}+\mathcal{K}}{P+1} - \gamma\bigg] + B_0\beta\alpha\zeta\bigg(\frac{\mathcal{M}}{2Q+S+1} - \gamma\bigg).$$

Canceling $\alpha\beta\zeta\hat{\beta}B_0$ we have

$$0 = \bigg[\frac{\mathcal{M}+\mathcal{K}}{P+1} - \gamma\bigg] - \bigg(\frac{\mathcal{M}}{2Q+S+1} - \gamma\bigg).$$

Remembering that $2Q+S = P$, the denominators are equal, and we have $0 = \mathcal{K}$, contradicting the hypothesis that the N-condition holds. That contradiction shows that the two terms do not cancel. But by the assumption that all the a_i and s_i are fast roots, we have $2Q+S+1 \leq P+1$. Hence by (22), $\operatorname{Im}\mathbb{H} = O(w^{P+1})$. That proves part (i) of the lemma.

It follows that the leading terms of the two parts of (20) do *not* cancel, and the asymptotic form of (20) is a constant times w^{P+1}. The constant is $\alpha\beta\zeta B_0\hat{\alpha}\hat{\beta}\hat{\zeta}\,[\mathcal{M}/(2Q+S+1)-\gamma]$. The first term of (19) has the leading term

$$\alpha\beta\zeta\hat{\alpha}\hat{\beta}\hat{\zeta}B_0\left(\frac{\mathcal{M}}{2Q+S+1}-\gamma\right)w^{2Q+S+1}$$

and the second term has the leading term

$$\alpha\beta\zeta\hat{\alpha}\hat{\beta}\hat{\zeta}B_0\left(\frac{\mathcal{M}+\mathcal{K}}{P+1}-\gamma\right)w^{P+1}$$

If there is at least one fast b_i then the second term has a higher power of w than the first. But even if the two terms have the same power of w, we have

$$\operatorname{Im}\mathbb{H} = O(w^{2Q+S+1})$$

Hence if $\mathbb{H}$ is not constant, we must have $Q = S = 0$, since otherwise $\operatorname{Im}\mathbb{H}$ will take two signs in the upper half plane. Thus all the a_i and s_i are principal, i.e. go to zero as t^{γ}. That completes the proof of the lemma.

Lemma 7 *If H is not constant, and the N-condition holds, then at each α_i, either $\mathbb{B}/\mathbb{A}$ has a simple zero or a simple pole, or is analytic non-vanishing at α_i.*

Proof. Let q be one of the α_i. Then by Lemma 3, q is real. We note that $\int_0^q \mathbb{A}^2\mathbb{S}\,dw > 0$ since $\mathbb{A}^2\mathbb{S}$ is, up to multiplication by a power of t and a factor $(1+O(t)$, the derivative u_x of the minimal surface u along the x-axis, which is nonnegative since u takes the boundary monotonically, and can be zero only at the origin. The following analysis works for $q = \beta_i$ as well, but then we do not start out knowing $\int_0^q \mathbb{A}^2\mathbb{S}\,dw \neq 0$.

We will make an analysis near q similar to the analysis we made above near zero. Thus the role played by the fast roots above will be played now by the roots that converge to q when t goes to zero. We re-use the letters "P", "Q", and "S" with new meanings: Let Q be the number of α_j that are equal to q, let P be the number of β_j that are equal to q, and let S be the number of ζ_j equal to q. Let $\hat{\beta}$ be the product of $\alpha - \beta_j$ over the j such that $\beta_j \neq q$, (so $\beta \neq 0$); let $\hat{\alpha}$ be the product of $q - \alpha_j$ over the j such that $\alpha_j \neq q$; and let ζ be the product of $q - \zeta_j$ over the j such that $\zeta_j \neq q$. We consider the behavior of the terms in (19) for w near q. Let $w = q + \xi$. We have

$$\begin{aligned}
\int_0^w \mathbb{A}^2\mathbb{S}\,dw &= \int_0^q \mathbb{A}^2\mathbb{S}\,dw + \alpha^2\zeta\hat{\alpha}^2\hat{\zeta}\frac{\xi^{2Q+S+1}}{2Q+S+1} + O(\xi^{2Q+S+2}) \\
\frac{\mathbb{B}}{\mathbb{A}} &= \frac{B_0\beta\hat{\beta}}{\alpha\hat{\alpha}}\xi^{P-Q}(1+O(\xi))
\end{aligned}$$

The coefficient of ξ in the $1+O(\xi)$ term is $\sum 1/(q-\beta_j) - \sum 1/(q-\alpha_j)$ where the sum is over the roots not equal to q; but the exact form of the error term does not matter. We have

$$\begin{aligned}
\frac{\mathbb{B}}{\mathbb{A}}\int_0^w \mathbb{A}^2\,dw &= \frac{B_0\beta\hat{\beta}}{\alpha\hat{\alpha}}\xi^{P-Q}\Big[\int_0^q \mathbb{A}^2\mathbb{S}\,dw \\
&\quad +\alpha^2\zeta\hat{\alpha}^2\hat{\zeta}\frac{\xi^{2Q+S+1}}{2Q+S+1}+O(\xi^{2Q+S+2})\Big](1+O(\xi)) \\
&= \frac{B_0\beta\hat{\beta}}{\alpha\hat{\alpha}}\int_0^q \mathbb{A}^2\mathbb{S}\,dw\;\xi^{P-Q}+\frac{B_0\alpha\beta\zeta}{2Q+S+1}\xi^{P+Q+S+1}+O(\xi^{P-Q+1}) \\
\int_0^w \mathbb{A}\mathbb{B}\mathbb{S}\,dw &= \int_0^q \mathbb{A}\mathbb{B}\mathbb{S}\,dw+B_0\alpha\beta\zeta\hat{\alpha}\hat{\beta}\hat{\zeta}\,\frac{\xi^{P+Q+1}}{P+Q+S+1}+O(\xi^{P+Q+S+2})
\end{aligned}$$

The last two lines can be combined using (19) to yield the following

$$H(w) \;=\; c_1+c_2\xi^{P-Q}+O(\xi^{P-Q+1})$$

where

$$c_2=(2m+1)B_0\frac{\beta\hat{\beta}}{\alpha\hat{\alpha}}\int_0^q \mathbb{A}^2\mathbb{S}\,dw.$$

We proved in the first paragraph of this proof that the integral appearing in c_2 is not zero, and $\beta\hat{\beta}/(\alpha\hat{\alpha})$ is not zero either, since it is a quotient of products of nonzero numbers. Unless $|P-Q|\le 1$, i.e. $P=Q\pm 1$ or $P=Q$, the imaginary part of the leading term $c_2\xi^{P-Q}$ will not have one sign in the upper half-plane. Therefore, $P=Q\pm 1$ or $P=Q$. That completes the proof of the lemma.

9 What is true if $\mathbb{H}$ is constant

In this section we draw out some consequences of the assumption that $\mathbb{H}$ is constant.

Lemma 8 *If H is constant, then*

$$\frac{\mathbb{B}}{\mathbb{A}}=\frac{B_0\beta}{\alpha}\Big(\int \mathbb{A}^2\mathbb{S}\,dw\Big)^{\mathcal{K}/\mathcal{M}}$$

and $\mathbb{B}/\mathbb{A}$ is a polynomial.

Proof. Let $\mathbf{Q}$ be the rational function $\mathbb{B}/\mathbb{A}$. The formula for $\mathbb{H}$ becomes

$$\begin{aligned}
\mathbb{H} &= \mathcal{M}\frac{\mathbb{B}}{\mathbb{A}}\int \mathbb{A}^2\mathbb{S}\,dw-(\mathcal{M}+\mathcal{K})\int \mathbb{A}\mathbb{B}\mathbb{S}\,dw \\
&= \mathcal{M}\mathbf{Q}\int \mathbb{A}^2\mathbb{S}\,dw-(\mathcal{M}+\mathcal{K})\int \mathbb{A}^2\mathbf{Q}\mathbb{S}\,dw
\end{aligned}$$

Differentiating with respect to w we have

$$\begin{aligned} \mathbb{H}' = 0 &= \mathcal{M}\mathbf{Q}' \int \mathbb{A}^2\mathbb{S}\,dw + \mathcal{M}\mathbf{Q}\mathbb{A}^2\mathbb{S} - (\mathcal{M}+\mathcal{K})\mathbf{Q}\mathbb{A}^2\mathbb{S} \\ &= \mathcal{M}\mathbf{Q}' \int \mathbb{A}^2\mathbb{S}\,dw - \mathcal{K}\mathbb{A}^2\mathbb{S}\mathbf{Q} \end{aligned}$$

The equation has one solution

$$\mathbf{Q} = \left(\int \mathbb{A}^2\mathbb{S}\,dw\right)^{\mathcal{K}/\mathcal{M}}$$

Since it is a linear equation, every solution is a multiple of that one; but as w goes to infinity, $\mathbf{Q}$ must be asymptotic to $B_0(\beta/\alpha)w^k$, so that condition determines the solution uniquely. We have now proved

$$\frac{\mathbb{B}}{\mathbb{A}} = \frac{B_0\beta}{\alpha}\left(\int \mathbb{A}^2\mathbb{S}\,dw\right)^{\mathcal{K}/\mathcal{M}}.$$

Multiplying by $\mathbb{A}$ we have

$$\mathbb{B} = \frac{B_0\beta}{\alpha}\mathbb{A}\left(\int \mathbb{A}^2\mathbb{S}\,dw\right)^{\mathcal{K}/\mathcal{M}}.$$

This equation shows that $\mathbb{B}$ vanishes wherever $\mathbb{A}$ does, and to at least as great an order. Hence the rational function $\mathbb{B}/\mathbb{A}$ is actually a polynomial. That completes the proof of the lemma.

Lemma 9 *Assume that N goes to $(-1,0,0)$ on compact subsets of the w-plane away from the zeroes of $\mathbb{A}$. If $\mathbb{H}(w)$ is constant, then, for the a_i and b_i that go to zero as t^γ, and R, Q, and S are the numbers of b_i, a_i, and s_i going to zero strictly faster than t^γ (the "fast roots") then we have $R = (2Q+S+1)\mathcal{K}/\mathcal{M}+Q$. In particular $R > Q$.*

Proof. By the previous lemma we have

$$\frac{\mathbb{B}}{\mathbb{A}} = \frac{B_0\beta}{\alpha}\left(\int \mathbb{A}^2\mathbb{S}\,dw\right)^{\mathcal{K}/\mathcal{M}}$$

Asymptotically near the origin we have

$$\int \mathbb{A}^2\mathbb{S}\,dw = \frac{\alpha^2\zeta}{2Q+S+1}w^{2Q+S+1}(1+O(w))$$

and hence

$$\left(\int \mathbb{A}^2\mathbb{S}\,dw\right)^{\mathcal{K}/\mathcal{M}} = \left(\frac{\alpha^2\zeta}{2Q+S+1}\right)^{\mathcal{K}/\mathcal{M}} w^{(2Q+S+1)\mathcal{K}/\mathcal{M}}(1+O(w))$$

and hence

$$\begin{aligned}
\mathbb{B} &= \frac{B_0\beta}{\alpha}\left(\frac{\alpha^2\zeta}{2Q+S+1}\right)^{\mathcal{K}/\mathcal{M}} \mathbb{A}\, w^{(2Q+S+1)\mathcal{K}/\mathcal{M}}(1+O(w)) \\
&= cw^{(2Q+S+1)\mathcal{K}/\mathcal{M}+Q}(1+O(w)) \qquad \text{for some constant } c \neq 0 \\
B_0\beta w^R &= cw^{(2Q+S+1)\mathcal{K}/\mathcal{M}+Q}(1+O(w))
\end{aligned}$$

Equating the exponents of the leading terms on left and right we have

$$R = (2Q+S+1)\mathcal{K}/\mathcal{M} + q$$

as required in (vii). That completes the proof of the lemma.

The following lemma reverses the above calculations, giving a sufficient condition for $\mathbb{H}$ to be constant.

Lemma 10 *Suppose $\mathcal{K} = \mathcal{M}(p-q)$ and $\mathbb{B}/\mathbb{A} = c(\int_0^w \mathbb{A}^2\, dw)^{p-q}$ for some constant c. Then $\mathbb{H}$ is constant.*

Proof. We have

$$\begin{aligned}
\mathbb{H} &= \mathcal{M}\frac{\mathbb{B}}{\mathbb{A}}\int_0^w \mathbb{A}^2 dw - (\mathcal{M}+\mathcal{K})\int_0^w \mathbb{A}\mathbb{B}\, dw \\
&= \mathcal{M}c\left(\int_0^w \mathbb{A}^2\, dw\right)^{p-q}\int_0^w \mathbb{A}^2\, dw - (\mathcal{M}+\mathcal{K})\int_0^w \mathbb{A}\mathbb{B}\, dw \\
&= \mathcal{M}c\left(\int_0^w \mathbb{A}^2 dw\right)^{p-q+1} - (\mathcal{M}+\mathcal{K})\int_0^w \mathbb{A}\mathbb{B}\, dw
\end{aligned}$$

Differentiating with respect to w, we have

$$\begin{aligned}
\frac{d\mathbb{H}}{dw} &= \mathcal{M}c(p-q+1)\left(\int \mathbb{A}^2\, dw\right)^{p-q}\mathbf{A}^2 - (\mathcal{M}+\mathcal{K})\mathbb{A}\mathbb{B} \\
&= \mathcal{M}(p-q+1)\left(\frac{\mathbb{B}}{\mathbb{A}}\right)\mathbb{A}^2 - (\mathcal{M}+\mathcal{K})\mathbb{A}\mathbb{B} \\
&= \mathcal{M}(p-q+1)\mathbb{A}\mathbb{B} - (\mathcal{M}+\mathcal{K})\mathbb{A}\mathbb{B} \\
&= (\mathcal{M}(p-q)+\mathcal{M})\mathbb{A}\mathbb{B} - (\mathcal{M}+\mathcal{K})\mathbb{A}\mathbb{B} \\
&= (\mathcal{M}(p-q)-\mathcal{K})\mathbb{A}\mathbb{B} \\
&= 0 \qquad \text{since } \mathcal{K} = \mathcal{M}(p-q) \text{ by hypothesis}
\end{aligned}$$

We have proved $d\mathbb{H}/dw = 0$; hence $\mathbb{H}$ is constant. That completes the proof of the lemma.

10 Gaussian area and the eigenfunction

We next draw some consequences from the calculations of the eigenfunction and the properties of H.

Lemma 11 *Suppose that on compact subsets of the w-plane away from the zeroes of $\mathbb{A}$, N converges to $(-1,0,0)$ as t goes to zero. Then it is not the case that both points $(1,0,0)$ and $(-1,0,0)$ (the "east pole" and "west pole") can be covered by N in the vicinity of one α_i.*

Proof. Fix a disk D_i centered at α_i and excluding all other α_i. Consider a point ξ in D_i at which the normal takes on $(1,0,0)$ or $(-1,0,0)$. We will show that if one of these is taken on in D_i, the other cannot be. At ξ we have ${}^1N = \pm 1$, and ${}^2N = {}^3N = 0$. The formula (17) calculated above for the eigenfunction ϕ in terms of N (before eliminating N in favor of $\tilde{\mathbb{A}}$ and $\tilde{\mathbb{B}}$) shows that at such a point ξ we have

$$\begin{aligned} \frac{2}{t^{\mathcal{M}-1}}\phi &= {}^1N\,\mathrm{Re}\left(\mathcal{M}\int_0^{\xi}\tilde{\mathbb{A}}^2\,\tilde{\mathbb{S}}\,dw-\tilde{\mathbb{A}}^2(\xi)\,\tilde{\mathbb{S}}(\xi)\,\xi+O(t)\right)\\ &= {}^1N\,c(1+O(\xi-\alpha_i))+O(t) \end{aligned}$$

where c is the constant $\mathcal{M}\int_0^{\alpha_i}\mathbb{A}^2\,\mathbb{S}\,d\xi$. The constant c is nonzero since (i) α_i is real, by Lemma 3, and and (ii) the integrand $\mathbb{A}^2\,\mathbb{S}$ is nonnegative on the real axis, because u takes the boundary monotonically and $u_x = t^{\mathcal{M}+2d+f}\mathbb{A}^2\,\mathbb{S}(1+O(t))$, and (iii) $\alpha_i \neq 0$, since a_i goes to zero as t^{γ}.

As t goes to zero, ξ will converge to α_i, since by assumption outside every disk centered at α_i, N converges to $(0,0,-1)$. (We do not require that ξ depend smoothly on t; a sequence of values of t going to zero will suffice, or even a single choice of t depending appropriately on the constants hidden in the $\mathbf{O}(t)$ and $O(w)$ terms.) Therefore the $O(\xi-\alpha_i)$ term goes to zero too. For small enough t, we see that the sign of $\phi(w)$ is equal to the sign of ${}^1N$ if $\alpha > 0$, or is opposite to the sign of ${}^1N$ if $\alpha < 0$. In particular, for all such points in the vicinity of a fixed α_i, the sign of ${}^1N$ is the same, since the sign of ϕ is the same in the entire upper half plane. That completes the proof.

The significance of this lemma is that the Gauss map, considered as a covering map, covers at most one hemisphere (plus $O(t)$), since if it could be extended across the narrow strip near the YZ plane where the image of the boundary must lie, it would reach the other of the two points in question. Of course it covers a full sphere (within $O(t)$) in a small disk around $\mathbf{a}_i$, since the normal takes on the north pole at each $\mathbf{a}_i(t)$, but it could conceivably take all 4π of its Gaussian area in the lower half plane, rather than one hemisphere in each half plane.

We say that α_i *contributes Gaussian area* if, in the vicinity of α_i, the normal restricted to a small upper half-disk covers an area at least $2\pi - O(t)$ on the sphere. To make this definition precise, let D_ρ be a disk of radius ρ about α_i in the ξ-plane, and let D_ρ^+ be its intersection with the upper half plane. Then α_i contributes Gaussian area if, for every $\rho > 0$, $N(D_\rho^+)$ has area at least $2\pi - O(t)$ for sufficiently small t.

Since the normal N is a projection mapping (away from its ramification points), and since the image of the boundary is confined to within $O(t)$ of the YZ-plane, as soon as N takes on a value more than $O(t)$ from the YZ-plane (in D_t^+) then α_i contributes Gaussian area. In particular, if N takes on the "east pole" $(1,0,0)$ or the "west pole" $(-1,0,0)$ in D_t^+ for every $t > 0$, then α_i contributes Gaussian area.

Lemma 12 *Suppose that on compact subsets of the w-plane away from the zeroes of $\mathbb{A}$, N converges to $(-1,0,0)$ as t goes to zero. If α_i contributes Gaussian area, then*

(i) There exist two points in the ξ-plane, say $eq_1(t)$ and $eq_2(t)$, both converging to α_i as t goes to zero, such that ${}^2N(eq_1(t)) = 1 + \mathbf{O}(t)$ and ${}^2N(eq_2(t)) = -1 + \mathbf{O}(t)$. In particular these two points have the same sign, since they both are near α_i.

(ii) $eq_i(t)$ can be found in such a way that $eq_i(t)$ has a limit as t goes to zero, and even depends analytically on a rational power of t. Therefore such expressions as $\tilde{\mathbb{A}}[\xi := eq_i(t)]$ are analytic in a rational power of t, and have limits as $t \to 0$.

Proof. The stereographic projection of the unit normal N is given by $t^{k\gamma}\tilde{\mathbb{B}}/\tilde{\mathbb{A}}$. The condition demanded in the lemma therefore has the geometric meaning that N should be horizontal at $eq(t)$. Pick a small number ρ. Let D be the disk of radius ρ about α_i; let D^+ be the part D lying in the upper half-plane. By hypothesis, α_i contributes Gaussian area, so it is not the case that $N(D^+)$ lies within $O(t)$ of the YZ plane. Then N does cover one of the points $(-1,0,0)$ and $(1,0,0)$. ${}^3N$ is $-1 + O(t^{2k})$ at the endpoints of the interval $[\alpha - \rho, \alpha + \rho]$, and indeed on the entire circle ∂D, by the hypothesis of the lemma.

Ad (i): In Lemma 11 we proved that the Gauss map in the vicinity of α_i covers at most one of the two possible hemispheres. By hypothesis it covers at least one of them, since it covers either $(-1,0,0)$ or $(1,0,0)$ and can be continued from there over either the half-space $X \geq O(t)$ or the half-space $X \leq O(t)$. The pre-image of the equator over D^+, that is, the set of points w in D^+ such that ${}^3N(w) = 0$, must lie in either the half-space $X \leq O(t)$ or the half-space $X \geq O(t)$; otherwise both points $(1,0,0)$ and $(-1,0,0)$ would be covered, contrary to Lemma 11. According to that lemma, at most one of those points is covered, and as remarked above, at least one of them is covered. Suppose that $(1,0,0)$ is covered (the argument is similar if $(-1,0,0)$ is covered). Since away from the boundary, N is a covering projection, we can find a path π in D^+ such that $N(\pi(t))$ lies on the equator, and ${}^1N(\pi(0)) = (1,0,0)$. This path can be continued in both directions (even through possible ramification points) until the image reaches $(-1,0,0)$, or until $\pi(t)$ reaches the boundary of D_t^+. If it does reach the boundary of D^+, it has to be on the real axis that it does so, since on the circular part of the boundary, $N(t) = O(t^{2k})$. If $\pi(t)$ reaches the boundary before $N(\pi(t))$ reaches within $O(t)$ of the XZ plane, then we have a contradiction, since the image under N of the boundary must lie within $O(t)$ of the XZ plane. If $N(\pi(t))$ reaches the XZ plane and passes more than $O(t)$ beyond the XZ plane before $\pi(t)$ reaches the boundary of D^+, then the path can be continued all the way to $(-1,0,0)$, since it cannot hit the boundary except when the image is within $O(t)$ of the XZ plane. But by Lemma 11, the path cannot be continued to $(-1,0,0)$. Hence $\pi(t)$ must reach the boundary of D^+ while $N(\pi(t))$ is within $O(t)$ of the XZ-plane. Then that value of $\pi(t)$ can be taken as $eq_i(t)$. Since the path on the sphere that goes along the equator can be continued in both directions, one end of π terminates in $eq_1(t)$ and the other end in $eq_2(t)$.

Ad (ii): The zero set of ${}^3N$ is defined by a quotient of functions analytic jointly in ξ and t. Its zero set therefore consists of a finite number of curves parametrizable as analytic functions of a rational power of t, by Lemma 1. That completes the proof.

Lemma 13 *Assume that* N *goes to* $(-1,0,0)$ *on compact subsets of the* w*-plane away from the zeroes of* $\mathbb{A}$. *If* α_i *contributes Gaussian area, then* $\mathbb{H}$ *is not constant.*

Proof. Suppose, for proof by contradiction, that $\mathbb{H}$ is constant. By Lemma 8 we have

$$\frac{\mathbb{B}}{\mathbb{A}} = \frac{B_0\beta}{\alpha}\left(\int_0^w \mathbb{A}^2\mathbb{S}\,dw\right)^{\mathcal{K}/\mathcal{M}} \tag{23}$$

By (10) we have

$${}^2N = \frac{(1+\mathbf{O}(t))2t^{2\mathcal{K}}\mathbb{A}\mathbb{B}}{\mathbb{A}^2+t^{2\mathcal{K}}\mathbb{B}^2}.$$

We now substitute $w := eq_i(t)$ into this equation, where $eq_1(t)$ and $eq_2(t)$ are the two points of Lemma 12. Then the two terms in the denominator are equal, and we obtain

$$\begin{aligned}
{}^2N[w := eq_i(t)] &= (1+\mathbf{O}(t))\frac{2t^{2k\gamma}\mathbb{A}\mathbb{B}}{2t^{2\mathcal{K}}\mathbb{B}^2} \\
&= (1+\mathbf{O}(t))\frac{\mathbb{A}}{\mathbb{B}}
\end{aligned}$$

Substituting for $\mathbb{A}/\mathbb{B}$ using (23) we have

$${}^2N[w := eq_i(t)] = \frac{1+\mathbf{O}(t)}{\left(B_0\int_0^{eq_i(t)}\mathbb{A}^2\mathbb{S}\,dw\right)^{\mathcal{K}/\mathcal{M}}}$$

The denominator is not zero, since $\mathbb{A}^2\mathbb{S} \geq 0$ because u takes the boundary monotonically, and $eq_i(t)$ converges to α_i. Since $eq_1(t)$ and $eq_2(t)$ both converge to α_i, they have the same sign. Therefore the right hand side of this equation has the same sign for $i=1$ and $i=2$. This contradicts Lemma 12, which says that the left-hand side takes values near 1 and -1 (in some order) for $i=1,2$ and t small but positive. That completes the proof.

11 Descending through the rings of roots

Recall that the roots a_i, b_i, and s_i fall into "rings", where each ring of roots goes to zero as t^{γ_n}, and $\gamma_1 > \gamma_2 \ldots$. For each ring we can consider the corresponding

w-plane, where $z = t^{\gamma_i} w$. In each ring we can consider the $\mathbb{H}$ for that ring. Let us abbreviate by the phrase "N-condition" the proposition that N goes to $(-1, 0, 0)$ on compact subsets of the w-plane apart from the zeros of $\mathbb{A}$. The N-condition depends on n, since the power of t used to define the w-plane for the n-th ring of roots is γ_n. If the n-th ring satisfies the N-condition, and $\mathbb{H}$ for ring n is constant, then by Lemma 9, there are more fast b_i's than there are fast a_i's, so the $n+1$-st ring also will satisfy the N-condition. This is the fundamental observation behind the following lemma:

Lemma 14 *(i) For some n, if $\gamma = \gamma_n$ and w and $\mathbb{H}$ are as defined above, then $\mathbb{H}$ is not constant.*

(ii) $\gamma = \gamma_1$, i.e. all roots belong to the first ring; all the a_i are principal and real, the s_i do not occur, and each a_i contributes 2π in Gaussian area.

(iii) At each α_i, $\mathbb{B}/\mathbb{A}$ has a simple pole or simple zero, or is analytic non-vanishing.

(iv) All the b_i are also principal.

Proof. According to the Gauss-Bonnet-Sasaki-Nitsche formula, some root a_i must contribute Gaussian area; in fact the total extra Gaussian area contributed in the upper half plane near origin must be $2m\pi$. Suppose, for proof by contradiction, that $\mathbb{H}$ is constant on all the rings of roots. Then, as sketched above and shown below in detail, the N-condition holds on all the rings. Then, by Lemma 13, no Gaussian area can be contributed. This contradiction will prove part (i).

It remains to check the details of the claim that the N-condition will hold on ring $n+1$ if it holds on ring n and $\mathbb{H}$ is constant on ring n. The N-condition is equivalent to $\tilde{\mathbb{B}}/\tilde{\mathbb{A}} = O(t)$. Let R_n and S_n be the number of roots that go to zero as or faster than γ_n, and let e_n be the sum of $n_i\gamma_i$ for $i < n$, where n_i is the number of b_i that go to zero as t_i^{γ}; and let d_n be the sum of $n_i\gamma_i$ for $i < n$, where n_i is the number of a_i that go to zero as t_i^{γ}. Then in the n-th ring we have

$$\begin{aligned} \frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}} &= t^{\mathcal{K}} \frac{B}{A} \\ &= t^{e_n - d_n + \gamma(R_n - Q_n)} \frac{B}{A} \end{aligned}$$

Assuming that the N-condition holds for ring n, we have

$$e_n - d_n + \gamma(R_n - Q_n) > 0.$$

When we pass to ring $n+1$, the roots that go to zero slower than t_{n+1}^{γ} include those that contributed $e_n - d_n$ at ring n, but also those that go to zero as t^{γ_n}. Those now contribute to $e_{n+1} - d_{n+1}$ the powers of t that they were contributing to $\mathbb{B}/\mathbb{A}$, making no net change in the power of t in $\mathbb{B}/\mathbb{A}$. The remaining roots, the roots that go to zero faster than t^{γ_n}, now contribute a factor $t^{\gamma_{n+1}}$ instead of the former t^{γ_n}. Therefore the N-condition will hold at ring $n+1$ if there are more of the b_i going to zero faster than t^{γ_n} than there are a_i going to zero faster than t^{γ_n}. But that has been proved in Lemma 9. That completes the proof of part (i).

Ad (ii). According to the Gauss-Bonnet-Sasaki-Nitsche formula, a total of $2\pi m$ in extra Gaussian area must be contributed by the a_i, of which there are $2m - N$. According to Lemma 11, in the vicinity of each α_i at most one hemisphere is covered. Since in a small disk about α_i the number of sheets over the north pole in the covering of the sphere by N is equal to the number of $\mathbf{a}_j$ that converge to that α_i, the Gaussian area contributed in the vicinity of α_i is 2π times the number of $\mathbf{a}_j$ that converge to that α_i. The $\mathbf{a}_i$ that go to zero slower than t^γ contribute nothing, by Lemma 13, since on those rings $\mathbb{H}$ is constant and the N-condition holds. By Lemma 6, part (ii), no a_i goes to zero faster than t^γ. Hence all the Gaussian area is contributed on the first ring of roots where $\mathbb{H}$ is not constant, and it is 2π times the number of a_i in that ring. But the total must be $2\pi m$, so there must be $2m$ of the a_i. Hence $N = 0$, i.e. the s_i do not occur. Moreover, by Lemma 6, part (i), in each ring there is at least one principal a_i or s_i; but we have just proved there are no s_i, so in each ring there is a principal a_i. Then the ring where $\mathbb{H}$ is not constant must be the first ring, since all $2m$ of the a_i occur in that ring. Moreover, by lemma 3, the α_i are real. That completes the proof of part (ii).

Ad (iii). This follows from Lemma 7 since $\mathbb{H}$ is not constant.

Ad (iv). If some of the b_i are "fast" roots, i.e. go to zero faster than t^γ, then the next ring contains some b_i; but by parts (i) and (ii), it contains no a_i or s_i. Since there are then more fast b_i's than a_i's, the next ring would also satisfy the N-condition. But then the fact that there are no a_i or s_i in the next ring contradicts Lemma 6, part (i). That completes the proof of the lemma.

Remark. Tony Tromba asked whether our proof requires an appeal to external references to know that the branch point is not a false branch point. It does not, and the key reason why not is our application of the Gauss-Bonnet-Sasaki-Nitsche formula. The surfaces u^t for $t > 0$ must contribute Gaussian area. Therefore $\mathbb{H}$ is not constant, as we have shown. That means that the surfaces u^t are not simply "sliding" over the same geometrical surface as t becomes positive; there is a normal component, with *some* power of t, to the motion. In the next section we show that $k/(2m+1)$ is an integer, which would be the case for a false branch point, but a minimal surface with a false branch point would have to be a function of z^{2m+1}, i.e., *every* term would have to have a power divisible by $2m + 1$, whereas intuitively it is the failure of this condition that makes $\mathbb{H}$ not constant, although we do not have a direct proof of that.

12 $k/(2m + 1)$ is an integer

In this section, we will deduce the result in the section title by analyzing the leading terms (leading power of t and its coefficient as a function of w) of the imaginary parts of $\tilde{\mathbb{A}}$, $\tilde{\mathbb{B}}$, and $\tilde{\mathbb{A}}\tilde{\mathbb{B}}$. Specifically, we are referring to the imaginary parts of these functions *on the boundary*, which may or may not have the same power of t as the same functions considered on the plane. Our plan for this analysis is simply to compare the expressions given for the surface on the boundary by (a) the analytic parametrization of the boundary and (b) the Weierstrass representation.

We begin with an elementary lemma.

Lemma 15 *Let f be meromorphic. For real w we have*

$$\mathrm{Im}\,(f_w) = 2\mathrm{Re}\,((\mathrm{Im}\, f)_w).$$

In particular if f is real for real w then $\mathrm{Im}\,(f_w) = 0$.

Proof. Let $f = U + iV$, with U and V real, and let $w = u + iv$. Then $f_w = U_w + iV_w = (U_u - iU_v)/2 + i(V_u - iV_v)/2$, so $\mathrm{Im}\,(f_w) = (-U_v + V_u)/2 = V_u$ by the Cauchy-Riemann equations. On the other hand $(\mathrm{Im}\, f)_w = V_w = (V_u - iV_v)/2$ and the real part of this is $V_u/2$. but this is $(\mathrm{Im}\, f)_w$ for real w. That completes the proof of the lemma.

It will shorten many of our formulas to use the following definition:

$$\sigma := \int_0^w \tilde{\mathbb{A}}^2\, dw \tag{24}$$

For example, the definition of H, given in (18) becomes

$$\mathbb{H}(w) = \frac{\mathbb{B}}{\mathbb{A}}\,(2m+1)\sigma - (2m+k+1)\int_0^w \mathbb{A}\mathbb{B}\, dw \tag{25}$$

Note that σ is defined using $\tilde{\mathbb{A}}$, not $\mathbb{A}$, so it depends on t and it has an imaginary part on the boundary. We claim that the leading powers of t in $\mathrm{Im}\,\tilde{\mathbb{A}}$ and $\mathrm{Im}\,\sigma$ are the same. To see this we differentiate the definition of σ with respect to w, to obtain

$$\sigma_w = \tilde{\mathbb{A}}^2$$

Taking the imaginary part we have $\mathrm{Im}\,(\sigma_w) = 2\tilde{\mathbb{A}}\mathrm{Im}\,\tilde{\mathbb{A}} = 2\mathbb{A}\mathrm{Im}\,\tilde{\mathbb{A}}(1 + O(t))$. Hence σ_w and $\tilde{\mathbb{A}}$ have the same leading power of t. Integrating with respect to w does not change the leading power of t, so σ has that same leading power, as claimed.

The main task in this section is to prove something about the leading powers of t in $\tilde{\mathbb{A}}$, $\tilde{\mathbb{B}}$, and $\tilde{\mathbb{A}}\tilde{\mathbb{B}}$. Eventually we want to characterize them in terms of the geometry of the boundary curve Γ. The following lemma gives a preliminary result.

Lemma 16 *Suppose that* $\mathrm{Im}\,\sigma$ *and* $\mathrm{Im}\,(\tilde{\mathbb{A}}\tilde{\mathbb{B}})$ *are not identically zero (on the boundary) for positive t. Then the leading power of t in* $\mathrm{Im}\,\tilde{\mathbb{A}}\tilde{\mathbb{B}}$ *is equal to the leading power of t in* $\mathrm{Im}\,\tilde{\mathbb{A}}$.

Proof. Whether or not $k/(2m+1)$ is an integer, we have, in a neighborhood of infinity,

$$\sigma^{k/(2m+1)} \quad = \quad \left(w^{2m+1}(1 + O(w^{2m}) + O(t))\right)^{k/(2m+1)}$$

$$= \quad w^k\left(1+O(w^{2m})+O(t)\right)^{k/(2m+1)}$$

The $O(t)$ term represents terms that may contain higher powers of w, but also contain at least one power of t. For fixed w, these terms go to zero with t. Thus the order of convergence implied is, holding w fixed, t goes to zero first. This expression is analytic in $1/w$ and t (for example because it can be expanded using the binomial series), so we have

$$\sigma^{k/(2m+1)} \quad = \quad w^k(1+O(1/w)+O(t))$$

where the $O(1/w)$ term is analytic in $1/w$ and t. Therefore the reciprocal of this expression has the form

$$\sigma^{-k/(2m+1)} \quad = \quad w^{-k}(1+O(1/w)+O(t))$$

Since also $\tilde{\mathbb{B}}/\tilde{\mathbb{A}} = w^k(1+O(1/w)+O(t))$, we have

$$\begin{aligned}\frac{\tilde{\mathbb{B}}/\tilde{\mathbb{A}}}{\sigma^{k/(2m+1)}} &= 1+O(1/w)+O(t) \\ &= 1+cw^{-J}+O(w^{-J-1})+O(t) \qquad (26)\end{aligned}$$

for some $J \geq 1$ and constant $c \neq 0$. Letting t go to zero we have

$$\frac{\mathbb{B}/\mathbb{A}}{\sigma^{k/(2m+1)}} \quad = \quad 1+cw^{-J}+O(w^{-J-1}) \qquad (27)$$

If $k/(2m+1)$ is an integer then we are dealing with a rational function, and we have $J \leq k$, since the worst case (largest J) is when all the coefficients in numerator and denominator are identical except the constant term. (Then $J = k$ and the constant c is the difference of the constant terms.) But if $k/(2m+1)$ is not an integer, we have no idea how large J might be.

Now we return to the main equation for $\mathbb{H}$, given above as (25):

$$\tilde{\mathbb{H}} = \frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}(2m+1)\sigma - (2m+k+1)\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\,dw$$

Taking the imaginary part we have

$$\operatorname{Im}\tilde{\mathbb{H}} \quad = \quad \operatorname{Im}\left(\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}(2m+1)\sigma\right) - \operatorname{Im}\left((2m+k+1)\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\,dw\right)$$

We can rewrite (26) as

$$\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}} = \sigma^{k/(2m+1)}(1+cw^{-J}+O(t)).$$

Putting this value into the previous equation we find

$$\begin{aligned}\operatorname{Im}\tilde{\mathbb{H}} \quad = \quad & \operatorname{Im}\left(\sigma^{k/(2m+1)}(1+cw^{-J}+O(t))(2m+1)\sigma\right) \\ & -\operatorname{Im}\left((2m+k+1)\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\,dw\right)\end{aligned}$$

Collecting powers of σ we have

$$\begin{aligned}\operatorname{Im}\tilde{\mathbb{H}} &= \operatorname{Im}\Big\{\sigma^{(2m+k+1)/(2m+1)}(2m+1)(1+cw^{-J}+O(t)) \\ &\quad -(2m+k+1)\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\,dw\Big\} \qquad (28)\end{aligned}$$

The imaginary part of $\sigma^{(2m+k+1)/(2m+1)}$ has the same leading power of t as the imaginary part of σ, as we shall now prove: Since $(\operatorname{Im}\sigma)/(\operatorname{Re}\sigma) = O(t)$ we have

$$\begin{aligned}\sigma^{(2m+k+1)/(2m+1)} &= \Big(\operatorname{Re}\sigma + i\operatorname{Im}\sigma\Big)^{(2m+k+1)/(2m+1)} \\ &= \Big((\operatorname{Re}\sigma)\Big(1+i\,\frac{\operatorname{Im}\sigma}{\operatorname{Re}\sigma}\Big)\Big)^{(2m+k+1)/(2m+1)} \\ &= (\operatorname{Re}\sigma)^{(2m+k+1)/(2m+1)}\Big(1+i\,\frac{\operatorname{Im}\sigma}{\operatorname{Re}\sigma}\Big)^{(2m+k+1)/(2m+1)}\end{aligned}$$

Expanding the power using the binomial series we have

$$\sigma^{(2m+k+1)/(2m+1)} = (\operatorname{Re}\sigma)^{(2m+k+1)/(2m+1)}\Big(1+i\,\frac{2m+k+1}{2m+1}\operatorname{Im}\sigma\Big)(1+O(t))$$

and taking the imaginary part we find

$$\operatorname{Im}\sigma^{(2m+k+1)/(2m+1)} = \frac{2m+k+1}{2m+1}\sigma^{(2m+k+1)/(2m+1)}\operatorname{Im}\sigma(1+O(t))$$

That proves the assertion that the imaginary part of $\sigma^{(2m+k+1)/(2m+1)}$ has the same leading power of t as the imaginary part of σ.

Now assume, for proof by contradiction, that the imaginary part of $\tilde{\mathbb{A}}\tilde{\mathbb{B}}$ has a higher power of t (on the boundary) than the imaginary part of σ (which is the same as the leading power of t in the imaginary part of $\tilde{\mathbb{A}}$). Therefore in (28), the last term is negligible, and outside of some large interval $[-R, R]$ centered at the origin, we have

$$\operatorname{Im}\tilde{\mathbb{H}} = \operatorname{Im}\Big\{\sigma^{(2m+k+1)/(2m+1)}(1+cw^{-J})(2m+1)\Big\}(1+O(t))$$

We can write the error term as a multiplicative factor $1+O(t)$ because the first term has the same leading power of t as $\operatorname{Im}\sigma$. The left-hand side must be zero for all t, since the eigenfunction vanishes on the boundary, but for sufficiently small (but positive) t, the right side is not zero, by the hypothesis that Im,σ is not identically zero.[5] This contradiction completes the proof that $\operatorname{Im}\tilde{\mathbb{A}}\tilde{\mathbb{B}}$ has its leading power of t less than or equal to the leading power of t in $\mathrm{Im}\,\sigma$. Now assume, for proof by contradiction, that strict inequality holds. Then in (28), the last term on the right dominates the first term on the right, and we have

---

[5] Note that these equations are all valid only on the boundary, so we are not considering the imaginary part of anything in the upper half plane here. In particular we are not appealing to the fact that $\operatorname{Im}\mathbb{H}$ has only one sign in the upper half plane.

$$\mathrm{Im}\,\tilde{\mathbb{H}} \quad = \quad -(2m+k+1)\mathrm{Im}\left\{\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\,dw\right\}(1+O(t))$$

But by the second hypothesis of the theorem, the right-hand side is not identically zero for positive t, contradiction. That completes the proof of the lemma.

We now state the main result of this section:

Lemma 17 *$k/(2m+1)$ is an integer.*

Proof. We will analyze the leading powers of t (on the boundary) in $\mathrm{Im}\,\tilde{\mathbb{A}}$, $\mathrm{Im}\,\tilde{\mathbb{B}}$, and $\mathrm{Im}\,\tilde{\mathbb{A}}\tilde{\mathbb{B}}$. In the following computations, we use $\cong$ instead of $=$ to indicate the omission of terms with higher powers of t than the leading powers of t in the terms that are explicitly shown. For example $\mathbb{A} \cong \tilde{\mathbb{A}}$; but we cannot write $\mathrm{Im}\,\mathbb{A} \cong \mathrm{Im}\,\tilde{\mathbb{A}}$ since the left side is zero. Since all these expressions are analytic in t and w, they can be differentiated, and as long as the first term is not constant, the results will still be $\cong$.

We recall that the boundary curve Γ is parametrized by a parameter τ (equal to the X-coordinate of the surface) so that

$$\Gamma' = \begin{bmatrix} 1 \\ C_2\tau^q(1+O(t)) \\ C_3\tau^p(1+O(t)) \end{bmatrix}$$

The point of using τ rather than arc length is that the first coordinate in this equation is exactly 1.

$$\begin{aligned}
u_t &= u_\tau \frac{d\tau}{dt} \\
&\cong u_\tau \frac{d}{dt}\left\{t^{(2m+1)\gamma}\int_0^w \mathbb{A}^2\,dw + t^{(2m+k+1)\gamma}\int_0^w \mathbb{B}^2\,dw\right\} \\
&\cong u_\tau\left(\frac{\partial}{\partial t} - \gamma t^{-1} w\frac{\partial}{\partial w}\right)\left\{t^{(2m+1)\gamma}\int_0^w \mathbb{A}^2\,dw + t^{(2m+k+1)\gamma}\int_0^w \mathbb{B}^2\,dw\right\} && \text{by (16)} \\
&\cong u_\tau \gamma t^{(2m+1)\gamma-1}\left\{(2m+1)\int_0^w \mathbb{A}^2\,dw - \mathbb{A}^2 w\right\} \\
&\cong \begin{bmatrix} 1 \\ C_2\tau^q \\ C_3\tau^p \end{bmatrix} \gamma t^{(2m+1)\gamma-1}\left\{(2m+1)\int_0^w \mathbb{A}^2\,dw - \mathbb{A}^2 w\right\}
\end{aligned}$$

On the other hand from the Weierstrass representation we have

$$u \quad \cong \quad t^{(2m+1)\gamma}\begin{bmatrix} \frac{1}{2}\mathrm{Re}\int_0^w \tilde{\mathbb{A}}^2 dw \\ \frac{1}{2}\mathrm{Im}\int_0^w \tilde{\mathbb{A}}^2\,dw \\ t^{k\gamma}\mathrm{Im}\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\,dw \end{bmatrix}$$

$$
\begin{aligned}
u_t &\cong \left(\frac{\partial}{\partial t} - \gamma t^{-1} w \frac{\partial}{\partial w}\right) u \\
&= \gamma t^{(2m+1)\gamma - 1} \left[\begin{array}{l} (2m+1)\int_0^w \mathbb{A}^2\, dw - \mathbb{A}^2\, w \\ \mathrm{Im}\left((2m+1)\int_0^w \tilde{\mathbb{A}}^2\, dw - \tilde{\mathbb{A}}^2\, w \right) \\ t^{k\gamma} (\mathrm{Im} \left((2m+k+1)\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\, dw - \tilde{\mathbb{A}}\tilde{\mathbb{B}}\, w \right) \end{array} \right]
\end{aligned}
$$

The first component ${}^1u_t$ gives identical results in both computations, but equating the second components in the two calculations, we find

$$
\begin{aligned}
& C_2 \gamma \tau^q t^{(2m+1)\gamma - 1} \left((2m+1) \int_0^w \mathbb{A}^2\, dw - \mathbb{A}^2\, w \right) \\
\cong\ & \gamma t^{(2m+1)\gamma - 1} \mathrm{Im} \left((2m+1) \int_0^w \tilde{\mathbb{A}}^2\, dw - \tilde{\mathbb{A}}^2\, w \right)
\end{aligned} \tag{29}
$$

Now

$$
\begin{aligned}
\tau^q &= \left(t^{(2m+1)\gamma} \int_0^w \mathbb{A}^2\, dw \right)^q (1 + O(t)) \\
&= t^{(2m+1)q\gamma} \left(\int_0^w \mathbb{A}^2\, dw \right)^q (1 + O(t))
\end{aligned}
$$

We simplify the appearance of that equation, and those to follow, by using σ as defined in (24), which we repeat here:

$$
\sigma := \int_0^w \tilde{\mathbb{A}}^2\, dw.
$$

Then the previous equation becomes

$$
\tau^q = t^{(2m+1)q\gamma} \sigma^q (1 + O(t))
$$

From (29) we have

$$
\begin{aligned}
& C_2 \gamma t^{(2m+1)q\gamma} t^{(2m+1)\gamma - 1} \sigma^q \left((2m+1)\sigma - \mathbb{A}^2\, w \right) \\
\cong\ & \gamma t^{(2m+1)\gamma - 1} \mathrm{Im} \left((2m+1)\sigma - \tilde{\mathbb{A}}^2\, w \right)
\end{aligned}
$$

Dividing both sides by $\gamma t^{(2m+1)\gamma - 1}$ we have

$$
C_2 t^{(2m+1)q\gamma} \sigma^q \left((2m+1)\sigma - \mathbb{A}^2\, w \right) \cong \mathrm{Im} \left((2m+1)\sigma - \tilde{\mathbb{A}}^2\, w \right) \tag{30}
$$

The coefficient on the left is not identically zero, since $(2m+1)\sigma - \mathbb{A}^2 w = 0$ if and only if $\mathbb{A} = w^m$, which cannot be the case since there is at least one principal a_i by Lemma 14. In detail: if $(2m+1)\sigma = \mathbb{A}^2 w$ then differentiating with respect to w, we have

$$\begin{aligned}
(2m+1)\mathbb{A}^2 &= 2w\mathbb{A}\mathbb{A}_w + \mathbb{A}^2 \\
2m\mathbb{A} &= 2w\mathbb{A}_w \\
\frac{\mathbb{A}_w}{\mathbb{A}} &= \frac{m}{w} \\
\log \mathbb{A} &= m \log w + c \\
&= \log c w^m \\
\mathbb{A} &= c w^m \\
&= w^m \qquad \text{since the leading coefficients on both sides are 1}
\end{aligned}$$

Having established that the coefficient on the left of (30) is not identically zero, we now have proved that the leading power of t on the left is the one shown, namely $t^{(2m+1)q\gamma}$. Hence that is also the leading power of t on the right, that is, in $\mathrm{Im}\,((2m+1)\sigma - \tilde{\mathbb{A}}^2 w)$. In particular, it is not the case that $\mathrm{Im}\,\sigma$ or $\mathrm{Im}\,\tilde{\mathbb{A}}$ is identically zero for positive t, since if either one is zero, so is the other one and then the right hand side of (30) would be zero, contradiction. Hence the first hypothesis of Lemma 16 is satisfied.

It would be nice to conclude that the imaginary parts of both σ and $\tilde{\mathbb{A}}^2$ also have leading term $t^{(2m+1)q\gamma}$, but there remains the possibility that the leading powers of t in the imaginary parts of σ and $\tilde{\mathbb{A}}^2$ are lower than $(2m+1)q\gamma$, but they cancel out. Hence, for now, we can only conclude that the leading powers of t in σ and in $\tilde{\mathbb{A}}$ (and hence $\tilde{\mathbb{A}}^2$) are the same, and are less than or equal to $(2m+1)q\gamma$.[6] We introduce the letter L for this power:

$$\mathrm{Im}\,\mathbb{A} \cong ct^L \qquad \text{for some } c$$

and we note that we have proved

$$L \le (2m+1)q\gamma \tag{31}$$

Similarly, comparing the two calculations of ${}^3u_t$, we find

$$\begin{aligned}
&C_3 t^{(2m+1)p\gamma} \sigma^p \left((2m+1)\sigma - \mathbb{A}^2\, w \right) \\
\cong\;& t^{k\gamma} \mathrm{Im} \left((2m+k+1) \int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\, dw - \tilde{\mathbb{A}}\tilde{\mathbb{B}}\, w \right)
\end{aligned}$$

As above, the coefficient on the left side is not identically zero as a function of w. Hence the right side is also not identically zero. It follows that $\mathrm{Im}\,(\tilde{\mathbb{A}}\tilde{\mathbb{B}})$ is not identically zero for positive t; hence the second hypothesis of Lemma 16 is also satisfied. Moreover, the leading power of t in $\operatorname{Im}(\tilde{\mathbb{A}}\tilde{\mathbb{B}})$ is less than or equal to $((2m+1)p-k)\gamma$.

---

[6] Note that the degree of the leading term in t of $\mathrm{Im}\,\mathbb{A}$, as a polynomial in w, could be much higher than $2m+1$; since everything was originally analytic in t and z, high powers of w must come with high powers of t, but there can also be lower powers of w that come with high powers of t. If $\mathrm{Im}\,\mathbb{A}$ really contains terms up to $w^{(2m+1)q}$, most of them must come from A_0, of which we have no direct knowledge, rather than from the $\mathbf{a}_i$, which might, for all we know, be *exactly* $\alpha_i t^\gamma$ with no higher powers of t involved.

Since both hypotheses of Lemma 16 have been verified, it follows from that lemma that $\operatorname{Im}\tilde{\mathbb{A}}\tilde{\mathbb{B}}$ has the same leading power of t as than $\operatorname{Im}\tilde{\mathbb{A}}$, namely t^L. In view of

$$\operatorname{Im}(\tilde{\mathbb{A}}\tilde{\mathbb{B}}) = \tilde{\mathbb{A}}\operatorname{Im}\tilde{\mathbb{B}} + \tilde{\mathbb{B}}\operatorname{Im}\tilde{\mathbb{A}}$$

and the fact that $\tilde{\mathbb{A}}$ and $\tilde{\mathbb{B}}$ are real when $t=0$, it follows that the leading power of t in $\operatorname{Im}\tilde{\mathbb{B}}$ is greater than or equal to L.

We now calculate the derivative of $\tilde{\mathbb{H}}$. Starting from

$$\tilde{\mathbb{H}} = (2m+1)\frac{\mathbb{B}}{\mathbb{A}}\int_0^w \tilde{\mathbb{A}}^2\,dw - (2m+k+1)\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\,dw$$

we find

$$\begin{aligned}
\tilde{\mathbb{H}}_w &= (2m+1)\frac{d}{dw}\left(\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\right)\int_0^w \tilde{\mathbb{A}}^2\,dw + (2m+1)\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\frac{d}{dw}\int_0^w \tilde{\mathbb{A}}^2\,dw \\
&\quad -(2m+k+1)\frac{d}{dw}\int_0^w \tilde{\mathbb{A}}\tilde{\mathbb{B}}\,dw \\
&= (2m+1)\left(\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\right)_w\int_0^w \tilde{\mathbb{A}}^2\,dw + (2m+1)\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\tilde{\mathbb{A}}^2 - (2m+k+1)\tilde{\mathbb{A}}\tilde{\mathbb{B}} \\
&= (2m+1)\left(\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\right)_w\int_0^w \tilde{\mathbb{A}}^2\,dw - k\tilde{\mathbb{A}}\tilde{\mathbb{B}}
\end{aligned}$$

This can be considered as a differential equation satisfied by $f = (\tilde{\mathbb{B}}/\tilde{\mathbb{A}})$:

$$\tilde{\mathbb{H}}_w = (2m+1)f'\int_0^w \tilde{\mathbb{A}}^2\,dw - k\tilde{\mathbb{A}}^2 f$$

or in terms of σ, the same equation reads

$$\tilde{\mathbb{H}}_w = (2m+1)f'\sigma - k\sigma_w f \tag{32}$$

We emphasize that in this section, we are only considering this equation on the boundary, that is, for real w. Since (even for $t>0$), the imaginary part of $\tilde{\mathbb{H}}$ is zero on the boundary, we also have $\operatorname{Im}(\tilde{\mathbb{H}}_w) = 0$ on the boundary, by Lemma 15.

Let $F(w)$ be the coefficient of the leading power of t, say t^J, in $\operatorname{Im}(\tilde{\mathbb{B}}/\tilde{\mathbb{A}})$. Then by definition of J and F we have

$$\operatorname{Im}\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}} = t^J F(w) + O(t^{J+1})$$

Then F is a quotient of real-analytic functions. We claim $J \ge L$. To prove this:

$$\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}} = \frac{\operatorname{Re}\tilde{\mathbb{B}} + \operatorname{Im}\tilde{\mathbb{B}}}{\operatorname{Re}\tilde{\mathbb{A}} + \operatorname{Im}\tilde{\mathbb{A}}}$$

$$
\begin{aligned}
&= (1+O(t))\frac{\tilde{\mathbb{B}}+\mathrm{Im}\,\tilde{\mathbb{B}}}{\mathbb{A}(1+\frac{\mathrm{Im}\,\tilde{\mathbb{A}})}{\mathbb{A}}} \\
&= (1+O(t))\frac{(\tilde{\mathbb{B}}+\mathrm{Im}\,\tilde{\mathbb{B}})(1-\frac{\mathrm{Im}\,\tilde{\mathbb{A}}}{\mathbb{A}})}{\mathbb{A}} \\
&= (1+O(t))\left\{\frac{\mathbb{B}}{\mathbb{A}}+\frac{\mathrm{Im}\,\tilde{\mathbb{B}}}{\mathbb{A}}-\frac{\tilde{\mathbb{B}}\,\mathrm{Im}\,\tilde{\mathbb{A}}}{\mathbb{A}^2}\right\} \\
\mathrm{Im}\left(\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\right) &= (1+O(t))\left\{\frac{\mathrm{Im}\,\tilde{\mathbb{B}}}{\mathbb{A}}-\frac{\mathbb{B}\,\mathrm{Im}\,\tilde{\mathbb{A}}}{\mathbb{A}^2}\right\}
\end{aligned}
$$

The leading power of t on the left is t^J, and on the right is either t^L or some larger power, which is possible if the terms on the right cancel. Hence

$$
J \geq L \tag{33}
$$

Since the $O(t^{J+1})$ terms above are analytic, they can be differentiated. Differentiating (33) we have

$$
\left(\mathrm{Im}\left(\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\right)\right)_w = t^J F_w + O(t^{J+1}) \tag{34}
$$

Taking the imaginary part of the differential equation (32) we have

$$
\begin{aligned}
\mathrm{Im}\,\tilde{\mathbb{H}}_w &= (2m+1)\mathrm{Im}\left(\left(\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\right)_w\right)\sigma+(2m+1)\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\mathrm{Im}\,\sigma-k\tilde{\mathbb{A}}^2\mathrm{Im}\,\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}-k\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\mathrm{Im}\,\tilde{\mathbb{A}}^2 \\
&= (2m+1)\mathrm{Re}\left(\left(\mathrm{Im}\left(\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\right)\right)_w\right)\sigma \\
&\quad +(2m+1)\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\mathrm{Im}\,\sigma-k\tilde{\mathbb{A}}^2\mathrm{Im}\,\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}-k\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\mathrm{Im}\,\tilde{\mathbb{A}}^2 \qquad \text{by Lemma 15} \\
&= (2m+1)\Big(t^J F_w\Big)\sigma+(2m+1)\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\mathrm{Im}\,\sigma \\
&\quad -k\tilde{\mathbb{A}}^2\mathrm{Im}\,\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}-k\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\mathrm{Im}\,\tilde{\mathbb{A}}^2+O(t^{J+1}) \text{ by (34)}
\end{aligned}
$$

$$
\begin{aligned}
&= (2m+1)\Big(t^J F_w\Big)\sigma+(2m+1)\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\mathrm{Im}\,\sigma \\
&\quad -k\tilde{\mathbb{A}}^2 t^J F-k\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\mathrm{Im}\,\tilde{\mathbb{A}}^2+O(t^{J+1}) \text{ by (33)}
\end{aligned}
$$

By definition of σ (see (24)), we have $\tilde{\mathbb{A}}^2 = \sigma_w$. Putting that into the previous equation and regrouping the terms, we have

$$
\begin{aligned}
0 &= (2m+1)\Big(t^J F_w\Big)\sigma-k\sigma_w t^J F \\
&\quad +\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\mathrm{Im}\left\{(2m+1)\sigma-k\mathrm{Im}\,\tilde{\mathbb{A}}^2\right\}+O(t^{J+1}) \text{ by (33)}
\end{aligned}
$$

On the left side we have exactly zero, even for positive t, by Lemma 15. We have

$$\operatorname{Im}\left\{(2m+1)\sigma - k\tilde{\mathbb{A}}^2\right\} \quad = \quad \operatorname{Im}\left\{(2m+1)\sigma - \tilde{\mathbb{A}}^2\right\} - \operatorname{Im}(k-1)\tilde{\mathbb{A}}^2$$

The first term has leading power $t^{(2m+1)q\gamma}$, and the second one has leading power t^L. We have $L \leq (2m+1)q\gamma$ by (31), so the leading power of t on the left side is exactly L. Define $G(t,w)$ by the following equation:

$$t^L G(t,w) := \operatorname{Im}\left\{(2m+1)\sigma - k\tilde{\mathbb{A}}^2\right\} \tag{35}$$

Then G is analytic in t and w, and is not identically zero when t is zero. Now we rewrite the previous equation, putting in $G(t,w)$:

$$0 \quad = \quad (2m+1)\left(t^J F_w\right)\sigma - k\sigma_w F t^J + t^L \frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}} G(t,w) + O(t^{J+1}) \text{ by (33)}$$

Dividing by t^J and letting t go to zero we have

$$0 \quad = \quad (2m+1)F_w\sigma - k\sigma_w F + t^{L-J}\frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}} G(t,w) + O(t)$$

The first two terms are analytic away from the α_i when $t=0$. Since G is not identically zero when $t=0$, we must have $L=J$. That is, the leading power of t in $\tilde{\mathbb{A}}$ and σ is the same as that in $\tilde{\mathbb{B}}/\tilde{\mathbb{A}}$. We can thus drop the factor t^{L-J}, since it is 1:

$$0 \quad = \quad (2m+1)F_w\sigma - k\sigma_w F + \frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}} G(t,w) + O(t)$$

Now divide the equation by $(2m+1)F\sigma$. We find

$$0 \quad = \quad \frac{F_w}{F} - \frac{k}{2m+1}\frac{\sigma_w}{\sigma} + \frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\frac{G(t,w)}{(2m+1)F\sigma} + O(t)$$

Define

$$R(t,w) := \frac{\tilde{\mathbb{B}}}{\tilde{\mathbb{A}}}\frac{G(t,w)}{(2m+1)F\sigma} + O(t) \tag{36}$$

where the $O(t)$ term is the one in the previous line, so we have

$$0 \quad = \quad \frac{F_w}{F} - \frac{k}{2m+1}\frac{\sigma_w}{\sigma} + R(t,w)$$

Then R is a quotient of real-analytic functions, which is analytic when $t=0$ and w is away from the α_i. Integrating the equation, we find

$$\begin{aligned}
\log F \quad &= \quad \frac{k}{2m+1}\log\sigma - \int_0^w R(t,w) + c \qquad \text{for some constant } c \\
&= \quad \log \sigma^{k/(2m+1)} - \int_0^w R(t,w) + c \\
&= \quad \log \frac{\sigma^{k/(2m+1)}}{c e^{\int_0^w R(t,w)}} \qquad \text{for some other constant } c
\end{aligned}$$

and raising e to both sides we find

$$F = \frac{\sigma^{k/(2m+1)}}{ce^{\int_0^w R(t,w)}}$$

The left side is real-analytic away from the $\mathbf{a}_i$. Since $\mathbb{H}$ is not constant, all the roots are principal, by Lemma 14. That is, none of the a_i converges to zero; hence F is analytic at the origin, and from (36) we see that R is also real-analytic at the origin when $t = 0$. It follows that $\sigma^{k/(2m+1)}$ is real-analytic at the origin. Since all the roots are principal, we have $\sigma = \int_0^w \tilde{\mathbb{A}}^2\, dw = \alpha w(1 + O(w) + O(t))$. Hence $\sigma^{k/(2m+1)}$ is analytic if and only if $k/(2m+1)$ is an integer. Therefore, $k/(2m+1)$ is an integer. That completes the proof of the lemma.

The following elementary algebraic lemma will be needed when the previous lemma is applied.

Lemma 18 *Let F and G be two different polynomials of degree at most k, each with constant coefficient 1. Then there exists an integer $J \le k$ and constant $c \neq 0$ such that for small ξ we have*

$$\frac{F(x)}{G(x)} = 1 + cx^J + O(x^{J+1})$$

Proof. Let

$$\begin{aligned} F(x) &= 1 + a_1 + \dots a_k x^k \\ G(x) &= 1 + b_1 + \dots b_k x^k \end{aligned}$$

and let J be the least number such that $a_J \neq b_J$. Then $J \le k$ since by hypothesis F and G are not equal. Define

$$h(x) := 1 + \dots + a_{J-1}x^{J-1}.$$

Then the degree of h is at most $J - 1$ (though it may be less) and we have $h(x) = 1 + O(x)$.

$$\begin{aligned} F(x) - h(x) &= a_J x^J + O(x^{J+1}) \\ G(x) - h(x) &= b_J x^J + O(x^{J+1}) \end{aligned}$$

Possibly $a_J = 0$ or $b_J = 0$ but not both, since $a_J \neq b_J$. Then

$$\begin{aligned} \frac{F(x)}{G(x)} &= \frac{h(x) + (F(x) - h(x))}{h(x) + (G(x) - h(x))} \\ &= \frac{1 + \frac{F(x)-h(x)}{h(x)}}{1 + \frac{G(x)-h(x)}{h(x)}} \\ &= \frac{1 + \frac{a_J x^J + O(x^{J+1})}{h(x)}}{1 + \frac{b_J x^J + O(x^{J+1})}{h(x)}} \\ &= \frac{1 + a_J x^J + O(x^{J+1})}{1 + b_J x^J + O(x^{J+1})} \qquad \text{since } h(x) = 1 + O(x) \end{aligned}$$

Now we can apply the identity

$$\frac{1}{1+\xi} = 1 - \xi + O(\xi^2)$$

to obtain

$$\begin{aligned} \frac{F(x)}{G(x)} &= (1 + a_J x^J + O(x^{J+1}))(1 - b_J x^J + O(x^{J+1})) + O((b_J x^J + O(x^{J+1}))^2) \\ &= 1 + (a_J - b_J)x^J + O(x^{J+1}) \end{aligned}$$

Since $a_J - b_J \neq 0$ we can define $c = a_J - b_J$. That completes the proof of the lemma.

13 Finiteness

We are going to prove a theorem limiting the number of relative minima of area with a given boundary. To state such a theorem precisely, we must fix a topology to use in defining "relative minimum." Once a topology is fixed, a relative minimum is a surface u such that there exists a neighborhood of u (in the chosen topology) such that no surface in that neighborhood has smaller area than u. As usual in Plateau's problem, we work with harmonic surfaces.

We wish to use the C^n topology. First consider whether it is a stronger theorem for smaller n or larger n. *A priori* it might be possible to decrease area by a C^0 variation but not by a C^n variation, but if area can be decreased by a C^n variation, that same variation counts as a C^0 variation. Therefore a C^0 relative minimum is automatically a C^n relative minimum, but not vice-versa. Thus if there were infinitely many C^0 relative minima, there would automatically be infinitely many C^n relative minima. Taking the contrapositive, if there cannot be infinitely many C^n relative minima, there cannot be infinitely many C^0 relative minima. Thus the theorem will be stronger, when we use the C^n topology for as large an n as possible.

Unfortunately, our result depends on the regularity result that a relative minimum cannot have an interior or boundary branch point, which at present is known only for the C^0 topology.[7] When and if this regularity result is proved for larger n, our proof will immediately apply to the C^n topology; but for now we can prove it for the C^0 case.

Theorem 3 *Let Γ be a real-analytic Jordan curve in R^3. Then Γ cannot bound infinitely many disk-type minimal surfaces which are relative minima of area in the C^0 topology. What is more, Γ cannot bound infinitely many immersed disk-type minimal surfaces whose least eigenvalue satisfies $\lambda_{\min} \geq 2$.*

---

[7] Interior regularity for relative minima is proved for the C^1 topology in [1], and Tromba has an unpublished proof of interior regularity for relative minima in the C^n topology, but boundary regularity is still open even for C^1.

Remark: As usual in Plateau's problem, we are counting two surfaces that differ only by a conformal reparametrization as the same surface.
Proof: First, observe that the second claim of the theorem implies the first, since relative minima of area are immersed, by [24] [no true interior branch points], [16] [no false branch points], and [15] [no true boundary branch points if the boundary is analytic], and satisfy $\lambda \geq 2$, since if $\lambda < 2$ then the second variation of area is negative.

In [2], the theorems of [7] about the structure of the set of minimal surfaces bounded by Γ are used to prove (in Theorem 5.1) that if Γ is a real-analytic Jordan curve in R^3, such that Γ cannot bound a one-parameter family of minimal surfaces u^t with u^t immersed for $t > 0$, and each u^t having least eigenvalue 2, then the conclusion of the present theorem holds. It therefore suffices to prove that Γ cannot bound such a family u^t.

In previous sections we have introduced notation which we continue to use. In Lemma 14 we have proved that $\mathbb{H}$ is not constant. Therefore by Lemma 14, the s_i do not occur, all the a_i are principal and real, and at each α_i, $\mathbb{H}$ has a simple zero or a simple pole. Until the previous section, we have examined $\mathbb{H}$ only in the neighborhood of particular points; and there, we examined its behavior at infinity only on the boundary. To complete the proof we will examine its behavior at infinity in the upper half plane.

As in the previous section, let $\sigma := \int_0^w \tilde{\mathbb{A}}^2 \, dw$. We have from (32) that $f = \mathbb{B}/\mathbb{A}$ is a solution of the differential equation

$$\frac{\mathbb{H}_w}{2m+1} = f'\sigma - \frac{k}{2m+1}\sigma_w f$$

Multiplying both sides by $\sigma^{k/(2m+1)-1}$ and dividing by $\sigma^{2k/(2m+1)}$ we have

$$\begin{aligned} \frac{\mathbb{H}_w}{(2m+1)\sigma^{k/(2m+1)+1}} &= \frac{f'\sigma^{k/(2m+1)} - \left(\sigma^{k/(2m+1)}\right)_w f}{\sigma^{2k/(2m+1)}} \\ &= \left(\frac{f}{\sigma^{k/(2m+1)}}\right)' \end{aligned}$$

Multiplying both sides by the denominator of the left side, we have

$$\mathbb{H}_w = (2m+1)\sigma^{k/(2m+1)+1}\left(\frac{f}{\sigma^{k/(2m+1)}}\right)'$$

Looking at the behavior for large w, we have

$$\begin{aligned} \sigma &= \int_0^w \mathbb{A}^2 \, dw = \frac{w^{2m+1}}{2m+1} + O(w^{2m}) \\ \sigma^{k/(2m+1)} &= c_3 w^k + O(w^{k-1}) \qquad \text{for some } c_3 \\ \sigma^{k/(2m+1)+1} &= c_3 w^{2m+k+1} + O(w^{2m+k}) \qquad (37) \\ f &= \mathbb{B}/\mathbb{A} = w^k + O(w^{k-1}) \end{aligned}$$

Hence $f/\sigma^{k/(2m+1)}$ has both numerator and denominator asymptotic to w^k. This fraction is not identically 1, since then $\mathbb{H}_w$ would be zero, and $\mathbb{H}$ would be constant, contradicting Lemma 14. By Lemma 17, $k/(2m+1)$ is an integer. Whether or not it is an integer, we have

$$\frac{f}{\sigma^{k/(2m+1)}} = 1 + cw^{-J} + O(w^{-J-1}) \tag{38}$$

for some $J \geq 1$ and constant c. But since $k/(2m+1)$ is an integer, we are dealing with a rational function, whose numerator and denominator are polynomials of degree k. In that case we have $J \leq k$, since the worst case (largest J) is when the numerator and denominator have all their corresponding coefficients the same except the constant coefficient, and in that case $J = k$. For a more rigorous proof, divide numerator and denominator by w^k and then introduce $\xi = 1/w$; now we have polynomials in ξ of degree k, each having constant coefficient 1. By Lemma 18 there is some $J \leq k$ and some nonzero constant c such that the ratio of these polynomials is asymptotic to $c\xi^J$. That is, the quotient of the original polynomials is asymptotic as w goes to infinity to cw^{-J}. With the bound $J \leq k$ established, we continue the argument as follows:

$$\begin{aligned}
\mathbb{H}_w &= (2m+1)\sigma^{k/(2m+1)+1}\left(\frac{f}{\sigma^{k/(2m+1)}}\right)' \\
&= (2m+1)w^{2m+k+1}(1 + cw^{-J} + O(w^{-J-1}))' \qquad \text{by (37) and (38)} \\
&= (2m+1)w^{2m+k+1}(-Jcw^{-J-1} + O(w^{-J-2})) \\
&= -J(2m+1)cw^{2m+k-J} + O(w^{2m+k+J-1})
\end{aligned}$$

It is permissible to differentiate the big-O term because it represents functions analytic in w. Integrating, we have

$$\mathbb{H} = \frac{-J(2m+1)c}{2m+k+J+1}w^{2m+k-J+1} + O(w^{2m+k-J})$$

The big-O term can be integrated as shown because it represents functions analytic in w. Since $\operatorname{Im}\mathbb{H}$ must have one sign in some neighborhood of infinity, we must have $2m + k - J + 1 = \pm 1$. That is, $J = 2m + k + 1 \pm -1$. Since $k/(2m+1)$ is an integer, and hence $J \leq k$ as shown above, subtracting J from each side we have

$$\begin{aligned}
0 &= 2m + 1 + (k - J) \pm 1 \\
&\geq 2m + (k - J) \\
&\geq 2m \qquad \text{since } k - J \geq 0 \\
&\geq 1
\end{aligned}$$

where we have used Lemma 14 to infer that $m \geq 1$. This contradiction completes the proof.

14 Open Problems and Conjectures

The theorem proved here could possibly be generalized by weakening any of these hypotheses: (i) that the boundary is real-analytic; (ii) that the surfaces considered are of the topological type of the disk; or (iii) that the surfaces considered are relative minima of area. All questions formed by weakening one or more of these hypotheses are open. These questions are all "equal conclusion" strengthenings of the theorem, formed by weakening the hypotheses. We will comment on these questions one by one below.

There is also an interesting "stronger conclusion" version of the theorem, in which we ask for an explicit bound on the number of relative minima as a function of the geometry of the boundary curve Γ. Our ignorance is scandalous: Not counting curves for which uniqueness is known, there is only one Jordan curve Γ known for which anyone can give an explicit bound on the number of solutions of Plateau's problem (of disk type) bounded by Γ. That curve is "Enneper's wire" Γ_r (the image of the circle of radius r in the usual parametrization of Enneper's surface), for values of r slightly larger than 1, and the bound is three. This bound follows from the theorem of Ruchert [25] that uniqueness holds up to $r = 1$ and the theorem of [6] characterizing the space of minimal surfaces bounded by Γ_r in the vicinity of Enneper's surface for $r = 1$. The following conjecture, for example, seems presently out of reach: if the real-analytic Jordan curve Γ has total curvature less than or equal to 6π, then it bounds at most two relative minima of area (of disk type). We do know that there can be no such estimate for the number of minimal surfaces (minimizing or not), since Böhme proved in [8] that for any $\epsilon > 0$ and any n, there is a real-analytic Jordan curve with total curvature less than $4\pi + \epsilon$ bounding more than n minimal surfaces. But these are branched surfaces, not relative minima.

Since Enneper's wire bounds two relative minima when the total curvature exceeds 4π, we could expect to string together sequences of Enneper's wires with small bridges connecting them, obtaining for each n a Jordan curve with total curvature $4n\pi + \epsilon$ bounding (at least) 2^n relative minima (as well as some unstable minimal surfaces). Since we seem to have to increase the total boundary curvature to get many solutions of Plateau's problem, we might hope that for some explicitly given function F, we might have, for all real-analytic Jordan curves Γ of total curvature less than $2\pi n$, there are at most $F(n)$ relative minima of area. As far as we know, this might be true with $F(n) = 2^{n-1}$. Thus: (when $n = 2$) curvature less than 4π implies uniqueness (which is known); when $n = 3$, curvature less than 6π implies at most two relative minima (the conjecture mentioned above); when $n = 4$, curvature less than 8π implies at most 4 relative minima (also, of course, not known).

Next we take up the possible "equal conclusion" versions. First, what about weakening the smoothness of the boundary to C^n? Then we have some troubles about the boundary branch point representation, but perhaps that can be solved adequately. A more pressing difficulty is that we then cannot control the dependence of the zeroes, poles, and ramification points of the Gauss map N on the parameter t. There is also a third difficulty: even the proof of Tomi's theorem that there can't be infinitely many absolute minima does not go through for C^∞ boundaries, because the regularity result that a minimum of area cannot have a boundary branch point is still an open problem for C^∞ boundaries.

Second, what about generalizing the theorem to other topological types of surfaces or boundaries? Frank Morgan has given an example [20] of a boundary consisting of four circles, such that it bounds a continuum of unstable minimal surfaces of arbitrarily high genus. That still leaves open the finiteness problem for relative minima. I conjecture that a system of real analytic Jordan curves cannot bound infinitely many relative minima of any fixed orientable topological type (that is, any fixed number of boundary components and Euler characteristic). For non-orientable surfaces, there is a problem in that the least eigenfunction no longer has one sign. Instead, it has just one nodal line. Thus even Tomi's argument that there cannot be a loop of minimal surfaces all of which are absolute minima of area fails to go through for surfaces of the type of the Möbius strip. That is the simplest open question for non-orientable surfaces: can there be a one-parameter family of absolute minima of surfaces of the type of the Möbius strip?

Incidentally, another paper of Morgan [19] gives an example of a curve in R^4 that bounds a continuum of absolute minima of disk type. Therefore we cannot generalize the theorem to higher dimensional spaces.

Third, what about dropping the requirement that the surfaces be relative minima of area? Then our methods say nothing at all, since they all depend on analysis of the least eigenvalue and its eigenfunction. The only known results (other than uniqueness theorems) along these lines are two "6π theorems", both of which assume that Γ is a real-analytic Jordan curve whose total curvature is less than or equal to 6π. In [4], it is proved that such a curve cannot bound infinitely many minimal surfaces (stable or unstable). Earlier, Nitsche proved that same conclusion (see §A29 of [22], p. 447) using the additional hypothesis that Γ bounds no minimal surface with branch points. At the time of Nitsche's proof, the forced Jacobi fields had not yet been discovered. In [4], the theory of the forced Jacobi fields is used.

Appendix: Dictionary of Notation

To assist the reader we provide in one place a summary of the notations used in more than one section of this paper. This summary is provided as a memory aid, not as a list of complete definitions. The symbols are more or less in order of introduction; there is no natural "lexicographical" order for such a variety of symbols.

X, Y, Z	coordinates in R^3
Γ	a real-analytic Jordan curve in R^3, tangent to the X-axis at the origin
$u = u^t$	a one-parameter family of minimal surfaces,

	analytic in t in some interval $[0, t_0]$
t	the parameter defining that one-parameter family
${}^2u$	the second component of the vector $u = ({}^1u, {}^2u, {}^3u)$.
$z = x + iy$	the parameter used to define u^t
	u^t is analytic in a closed upper half-disk in the z-plane
N	the unit normal to u^t
g	the stereographic projection of the Gauss map N
u_t	subscript denotes differentiation: $\frac{\partial u}{\partial t}$
$\lambda_{\min}$	least eigenvalue
ϕ	$u_t \cdot N$. If nonzero, this is an eigenfunction for eigenvalue 2.
n and χ	$\phi = t^n \chi$.
f	${}^1u_z - {}^2u_z$, occurs in the Weierstrass representation of u^t
$2m$	order of the boundary branch point of u (when $t = 0$).
s_i	branch points in the lower half plane for $t > 0$ that converge to 0 as $t \to 0$.
N	There are $2N$ of the s_i. Hopefully no confusion will arise with N the unit normal.
a_i	zeroes of f. The a_i depend analytically on t. There are $m - N$ of these.
b_i	zeroes of fg^2. The b_i depend analytically on t. There are $m + k - N$ of these.
γ_n	The n-th ring of roots consists of a_i, b_i, and s_i that go to zero as t^{γ_n}.
γ	One of the γ_n.
principal roots	those a_i, b_i and s_i whose leading term is t^{γ_1}
α_i	coefficient such that $a_i = \alpha_i t^\gamma + O(t^{\gamma+1})$
β_i	coefficient such that $b_i = \beta_i t^\gamma + O(t^{\gamma+1})$
ζ_i	coefficient such that $s_i = \zeta_i t^\gamma + O(t^{\gamma+1})$
	α_i, β_i, and ζ_i are used only when non-zero, i.e. for the roots that go to zero as t^γ.
α	product of all the $-\alpha_i$ over roots going to zero slower than t^γ
β	product of all the $-\beta_i$ over roots going to zero slower than t^γ
ζ	product of all the $-\zeta_i$ over roots going to zero slower than t^γ
A	$A = A_0 \prod_{i=1}^{m-N} (z - a_i)$ so $f = A^2$, and $A_0 = 1$ when $t = 0$
B	$B = B_0 \prod_{i=1}^{m+k-N} (z - b_i)$ so $fg^2 = B^2$
B_0	the constant in the previous line
S	$S = \prod_{i=1}^{2N} (z - s_i(t))$
C	complex constant, value of B_0 when $t = 0$
$w = z/t^\gamma$	a "blow-up" of the parameter domain
$O(t)$	refers to convergence on compact subsets of the w-plane away from the α_i
$\mathbf{O}(t)$	refers to convergence on compact subsets of the w-plane (possibly including the α_i)
$q \cong r$	when expressed in terms of w and t, the coefficients of the lowest power of t in q and r agree as functions of w.
	That is: $q = r(1 + \mathbf{O}(t)$.
$\mathbb{A}$	$\mathbb{A} = \prod_{i=1}^{m-N} (w - \alpha_i)$ so $A = t^{(m-N)\gamma} \mathbb{A}(1 + \mathbf{O}(t))$
	$\mathbb{A}$ does not depend on t
$\mathbb{B}$	$\mathbb{B} = C \prod_{i=1}^{m+k-N} (w - \beta_i)$ so $B = t^{(m+k-N)\gamma} \mathbb{B}(1 + \mathbf{O}(t))$
	$\mathbb{B}$ does not depend on t, and C turns out to be real.
$\tilde{\mathbb{A}}$	$A(z) = t^{(m-N)\gamma} \tilde{\mathbb{A}}(t, w)$ defines $\tilde{\mathbb{A}}$

$\tilde{\mathbb{B}}$	$B(z) = t^{(m+k-N)\gamma}\tilde{\mathbb{B}}(t,w)$ defines $\tilde{\mathbb{B}}$
$\tilde{\mathbb{S}}$	$S(z) = t^{2N\gamma}\tilde{\mathbb{S}}(t,w)$ defines $\tilde{\mathbb{S}}$
$\mathbf{a}_i$	$\mathbf{a}_i = a_i/t^\gamma$
$\mathbf{b}_i$	$\mathbf{b}_i = b_i/t^\gamma$
$\mathbf{s}_i$	$\mathbf{s}_i = s_i/t^\gamma$
$\tilde{\mathbb{A}}_0(t,w)$	$\tilde{\mathbb{A}}_0(t,w) = A_0(t, t^\gamma w)$
$\tilde{\mathbb{B}}_0(t,w)$	$\tilde{\mathbb{B}}_0(t,w) = B_0(t, t^\gamma w)$
χ	$\chi(w) = \frac{\phi(t^\gamma w)}{\gamma t^{(2m+k+1)\gamma-1}}$
	χ shows what happens to the eigenfunction in the w-plane.
χ^0	limit of χ when t goes to 0
$\mathbb{H}$	complex analytic function such that $\chi^0 = -\mathrm{Im}\,\mathbb{H}$
σ	abbreviation for $\int_0^w \tilde{\mathbb{A}}^2\,dw$ in the last two sections
τ	the parameter used to parametrize the boundary curve, equal to the X-coordinate

References

[1] Beeson, M., On interior branch points of minimal surfaces, Math Zeitschrift **171** (1980) 133-154.

[2] Beeson, M., Some results on finiteness in Plateau's problem, Part I, *Math. Zeitschrift* **175** (1980) 103-123.

[3] Beeson, M. Some results on finiteness in Plateau's problem, Part II, *Math. Zeitschrift* **181** (1982) 1-30.

[4] Beeson, M., The 6π theorem about minimal surfaces, *Pacific Journal of Mathematics* **117** No. 1, 1985.[8]

[5] Beeson, M., www.MichaelBeeson.com/research/minimalsurfaces/BranchPointAnimations

[6] Beeson, M., and Tromba, A. J., The cusp catastrophe of Thom in the bifurcation of minimal surfaces, *Manuscripta Mathematica* **46** (1984) 273-308.

[7] Böhme, R., Die Zusammenhangskomponenten der Lösungen analytischer Plateauprobleme. *Math. Zeitschrift* **133**, 31-40 (1973).

[8] Böhme, R., New results on the classical problem of Plateau on the existence of many solutions. Séminaire Bourbaki 34 année **579** (1981–82) 1–20.

[9] Courant, R., *Dirichlet's Principle, Conformal Mapping, and Minimal Surfaces*, Interscience, New York (1950), reprinted by Springer-Verlag (1977).

---

[8] See also a revised version available on the author's web site, in which a direct argument is given that removes any dependence on [3].

[10] Dierkes, U., Hildebrant, S., Küster, A., Wohlrab, O., *Minimal Surfaces I: Boundary Value Problems*, Springer-Verlag, Berlin/Heidelberg/New York (1992).

[11] Dierkes, U., Hildebrant, S., Küster, A., Wohlrab, O., *Minimal Surfaces II: Boundary Regularity*, Springer-Verlag, Berlin/Heidelberg/New York (1992).

[12] Evans, L., *Partial Differential Equations*, Graduate Studies in Mathematics, vol. 19, American Mathematical Society, Providence, R. I. (1998).

[13] Gilbarg, D., and Trudinger, N. S., *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin/Heidelberg/New York (1977).

[14] Gulliver, R. A minimal surface with an atypical boundary branch point, in: B. Lawson and K. Tenenblat (eds.), *Differential Geometry: a Symposium in honor of Manfredo P. do Carmo*, pp. 211-228, Longman, Harlow (1991).

[15] Gulliver, R., and Lesley, F., On boundary branch points of minimizing surfaces, *Arch. Rational Mech. Anal.* **52**, 20–25 (1973).

[16] Gulliver, R., Osserman, R., and Royden, H., A theory of branched immersions of surfaces, *Amer. J. Math.* **95**, 750–812 (1973).

[17] Kato, T., *Perturbation Theory for Linear Operators*, second edition. Springer-Verlag, Berlin/Heidelberg/New York (1980).

[18] Lewy, H. On the boundary behaviour of minimal surfaces, Proc. Natl. Acad. Sci. USA **37**, 102-110 (1951).

[19] Morgan, F., A smooth curve in R^4 bounding a continuum of area minimizing surfaces, *Duke Math. J.* **43** (1976), 867–870.

[20] Morgan, F., A smooth curve in R^3 bounding a continuum of minimal manifolds, *Arch. Rat. Mech. Anal.* **75** (1981), 193–197.

[21] Morrey, C., Jr., *Multiple Integrals in the Calculus of Variations*, Springer-Verlag, Berlin/Heidelberg/New York (1966).

[22] Nitsche, J. C. C. *Lectures on Minimal Surfaces, Volume 1*, Cambridge University Press, Cambridge (1988).

[23] Osserman, R., *A Survey of Minimal Surfaces*, Dover Publications, New York (1986).

[24] Osserman, R., A proof of the regularity everywhere of the classical solution of Plateau's problem, *Annals of Mathematics* **91**, 550-569 (1970).

[25] Ruchert, H., A uniqueness result for Enneper's minimal surface, Indiana Univ. Math. J., **30(3)**, 427–431 (1981).

[26] Tomi, F., On the local uniqueness of the problem of least area, Arch. Ration. Mech. Anal. **52**, 312-381 (1978).

[27] Tomi, F., On the finite solvability of Plateau's problem, Lecture Notes in Mathematics **597**, 679–695, Springer, Berlin/Heidelberg/New York (1977).

[28] Tomi, F., and Tromba, A. J., On the structure of the set of curves bounding minimal surfaces of prescribed degeneracy, J. für die reine und angewandte Mathematik **316** (1980) 31–43.

[29] Tromba, A. J., On the number of simply connected minimal surfaces spanning a curve, Mem. Am. Math. Soc. No. 194, **12** (1977).